

Continuous orbit equivalence, flow equivalence of Markov shifts and circle actions on Cuntz–Krieger algebras

Kengo Matsumoto
Department of Mathematics
Joetsu University of Education
Joetsu, 943-8512, Japan

Abstract

We will study circle actions on Cuntz–Krieger algebras trivially acting on its canonical maximal abelian C^* -subalgebras from the view points of continuous orbit equivalence of one-sided topological Markov shifts and flow equivalence of two-sided topological Markov shifts.

1 Introduction and Preliminaries

For an $N \times N$ irreducible matrix $A = [A(i, j)]_{i,j=1}^N$ with entries in $\{0, 1\}$, let us denote by X_A the shift space

$$X_A = \{(x_n)_{n \in \mathbb{N}} \in \{1, \dots, N\}^{\mathbb{N}} \mid A(x_n, x_{n+1}) = 1 \text{ for all } n \in \mathbb{N}\} \quad (1.1)$$

of the one-sided topological Markov shift for A . We assume that $1 < N \in \mathbb{N}$ and A is not any permutation matrix. Hence A satisfies condition (I) in the sense of Cuntz–Krieger [8]. We endow X_A with the relative topology of the product topology on $\{1, \dots, N\}^{\mathbb{N}}$, so that X_A is a compact Hausdorff space and homeomorphic to the Cantor discontinuum. Let us denote by σ_A the shift map on X_A which is defined by $\sigma_A((x_n)_{n \in \mathbb{N}}) = (x_{n+1})_{n \in \mathbb{N}}$. It is a continuous surjection on X_A . The topological dynamical system (X_A, σ_A) is called the one-sided topological Markov shift for A . The topological Markov shifts are often called shifts of finite type (cf. [10], [11]). The two-sided topological Markov shift is similarly defined as a topological dynamical system $(\bar{X}_A, \bar{\sigma}_A)$ on the set $\bar{X}_A$ of two-sided sequences $(x_n)_{n \in \mathbb{Z}}$ instead of one-sided sequences $(x_n)_{n \in \mathbb{N}}$ in (1.1) with the shift homeomorphism $\bar{\sigma}_A((x_n)_{n \in \mathbb{Z}}) = (x_{n+1})_{n \in \mathbb{Z}}$ on $\bar{X}_A$. J. Cuntz and W. Krieger have introduced a C^* -algebra $\mathcal{O}_A$ associated to topological Markov shift (X_A, σ_A) ([8]). The C^* -algebra is called the Cuntz–Krieger algebra, which is a universal unique C^* -algebra generated by partial isometries $S_1, \dots, S_N$ subject to the relations:

$$\sum_{j=1}^N S_j S_j^* = 1, \quad S_i^* S_i = \sum_{j=1}^N A(i, j) S_j S_j^*, \quad i = 1, \dots, N. \quad (1.2)$$

For $t \in \mathbb{R}/\mathbb{Z}$, the correspondence $S_i \rightarrow e^{2\pi\sqrt{-1}t} S_i$, $i = 1, \dots, N$ gives rise to an automorphism of $\mathcal{O}_A$ denoted by ρ_t^A . The automorphisms ρ_t^A , $t \in \mathbb{R}/\mathbb{Z}$ yield an action of the circle

group $\mathbb{R}/\mathbb{Z}$ on $\mathcal{O}_A$ called the gauge action. Let us denote by $\mathcal{D}_A$ the C^* -subalgebra of $\mathcal{O}_A$ generated by the projections of the form: $S_{i_1} \cdots S_{i_n} S_{i_n}^* \cdots S_{i_1}^*, i_1, \dots, i_n = 1, \dots, N$. The subalgebra $\mathcal{D}_A$ is canonically isomorphic to the commutative C^* -algebra $C(X_A)$ of the complex valued continuous functions on X_A by identifying the projection $S_{i_1} \cdots S_{i_n} S_{i_n}^* \cdots S_{i_1}^*$ with the characteristic function $\chi_{U_{i_1 \dots i_n}} \in C(X_A)$ of the cylinder set $U_{i_1 \dots i_n}$ for the word $i_1 \cdots i_n$. Cuntz and Krieger have proved that if one-sided topological Markov shifts (X_A, σ_A) and (X_B, σ_B) are topologically conjugate, then there exists an isomorphism $\Phi : \mathcal{O}_A \rightarrow \mathcal{O}_B$ such that $\Phi(\mathcal{D}_A) = \mathcal{D}_B$ and $\Phi \circ \rho_t^A = \rho_t^B \circ \Phi, t \in \mathbb{R}/\mathbb{Z}$ ([8, 2.17 Proposition]).

Continuous orbit equivalence of one-sided topological Markov shifts is a weaker equivalence relation than one-sided topological conjugacy and gives rise to isomorphic Cuntz–Krieger algebras ([14]). Let A and B be irreducible square matrices with entries in $\{0, 1\}$. One-sided topological Markov shifts (X_A, σ_A) and (X_B, σ_B) are said to be continuously orbit equivalent if there exist a homeomorphism $h : X_A \rightarrow X_B$ and continuous functions $k_1, l_1 : X_A \rightarrow \mathbb{Z}_+, k_2, l_2 : X_B \rightarrow \mathbb{Z}_+$ such that

$$\sigma_B^{k_1(x)}(h(\sigma_A(x))) = \sigma_B^{l_1(x)}(h(x)) \quad \text{for } x \in X_A, \quad (1.3)$$

$$\sigma_A^{k_2(y)}(h^{-1}(\sigma_B(y))) = \sigma_A^{l_2(y)}(h^{-1}(y)) \quad \text{for } y \in X_B. \quad (1.4)$$

In [17], it has been proved that the following three assertions are equivalent (cf. [14], [15], [19], [20]):

- (i) (X_A, σ_A) and (X_B, σ_B) are continuously orbit equivalent.
- (ii) There exists an isomorphism $\Phi : \mathcal{O}_A \rightarrow \mathcal{O}_B$ such that $\Phi(\mathcal{D}_A) = \mathcal{D}_B$.
- (iii) $\mathcal{O}_A$ and $\mathcal{O}_B$ are isomorphic and $\text{sgn}(\det(\text{id} - A)) = \text{sgn}(\det(\text{id} - B))$.

Let g be a strictly positive continuous function on $\bar{X}_A$. Denote by $\bar{X}_A^g$ the compact Hausdorff space obtained from

$$\{(\bar{x}, r) \in \bar{X}_A \times \mathbb{R} \mid \bar{x} \in \bar{X}_A, 0 \leq r \leq g(\bar{x})\}$$

by identifying $(\bar{x}, g(\bar{x}))$ and $(\bar{\sigma}_A(\bar{x}), 0)$ for each $\bar{x} \in \bar{X}_A$. Let $\bar{\sigma}_{A,t}^g, t \in \mathbb{R}$ be the flow on $\bar{X}_A^g$ defined by

$$\bar{\sigma}_{A,t}^g([\bar{x}, r]) = [\bar{x}, r + t] \quad \text{for } [\bar{x}, r] \in \bar{X}_A^g.$$

The dynamical system $(\bar{X}_A^g, \bar{\sigma}_A^g)$ is called the suspension flow of $(\bar{X}_A, \bar{\sigma}_A)$ by ceiling function g . Two-sided topological Markov shifts $(\bar{X}_A, \bar{\sigma}_A)$ and $(\bar{X}_B, \bar{\sigma}_B)$ are said to be flow equivalent if for a strictly positive continuous function g on $\bar{X}_A$ there exists a strictly positive continuous function $\tilde{g}$ on $\bar{X}_B$ such that $(\bar{X}_A^g, \bar{\sigma}_A^g)$ is topologically conjugate to $(\bar{X}_B^{\tilde{g}}, \bar{\sigma}_B^{\tilde{g}})$. Throughout the paper, denote by $\mathcal{K}$ the C^* -algebra of compact operators on the separable infinite dimensional Hilbert space $\ell^2(\mathbb{N})$ and $\mathcal{C}$ its maximal abelian C^* -subalgebra consisting of diagonal elements on $\ell^2(\mathbb{N})$. We have seen that the following three assertions are equivalent ([17], [9], [28] cf. [2], [8], [22]) :

- (i) $(\bar{X}_A, \bar{\sigma}_A)$ and $(\bar{X}_B, \bar{\sigma}_B)$ are flow equivalent.
- (ii) There exists an isomorphism $\bar{\Phi} : \mathcal{O}_A \otimes \mathcal{K} \rightarrow \mathcal{O}_B \otimes \mathcal{K}$ such that $\bar{\Phi}(\mathcal{D}_A \otimes \mathcal{C}) = \mathcal{D}_B \otimes \mathcal{C}$.

(iii) $\mathcal{O}_A \otimes \mathcal{K}$ and $\mathcal{O}_B \otimes \mathcal{K}$ are isomorphic and $\text{sgn}(\det(\text{id} - A)) = \text{sgn}(\det(\text{id} - B))$.

Hence the continuous orbit equivalence of one-sided topological Markov shifts is regarded as a one-sided counterpart of the flow equivalence of two-sided topological Markov shifts.

Let us denote by $(\bar{H}^A, \bar{H}_+^A)$ the ordered cohomology group for the two-sided topological Markov shift $(\bar{X}_A, \bar{\sigma}_A)$, which is defined as the quotient group of the ordered abelian group $C(\bar{X}_A, \mathbb{Z})$ of all $\mathbb{Z}$ -valued continuous functions on $\bar{X}_A$ by the subgroup $\{\xi - \xi \circ \bar{\sigma}_A \mid \xi \in C(\bar{X}_A, \mathbb{Z})\}$. The classes of nonnegative functions in $C(\bar{X}_A, \mathbb{Z})$ form the positive cone $\bar{H}_+^A$ (cf. [1], [23]). The ordered cohomology group (H^A, H_+^A) for the one-sided topological Markov shift (X_A, σ_A) is similarly defined ([17]). The latter ordered group (H^A, H_+^A) is naturally isomorphic to the former one $(\bar{H}^A, \bar{H}_+^A)$ ([17, Lemma 3.1]). In [1], Boyle–Handelman have proved that the ordered group $(\bar{H}^A, \bar{H}_+^A)$ is a complete invariant for flow equivalence of two-sided topological Markov shift $(\bar{X}_A, \bar{\sigma}_A)$. Suppose that (X_A, σ_A) and (X_B, σ_B) are continuously orbit equivalent via a homeomorphism $h : X_A \rightarrow X_B$ satisfying (1.3) and (1.4). The homomorphism $\Psi_h : C(X_B, \mathbb{Z}) \rightarrow C(X_A, \mathbb{Z})$ defined by

$$\Psi_h(f)(x) = \sum_{i=0}^{l_1(x)-1} f(\sigma_B^i(h(x))) - \sum_{j=0}^{k_1(x)-1} f(\sigma_B^j(h(\sigma_A(x)))) \quad (1.5)$$

for $f \in C(X_B, \mathbb{Z})$, $x \in X_A$ and its inverse $\Psi_{h^{-1}} : C(X_A, \mathbb{Z}) \rightarrow C(X_B, \mathbb{Z})$ give rise to isomorphisms of abelian groups between $C(X_A, \mathbb{Z})$ and $C(X_B, \mathbb{Z})$. They furthermore induce isomorphisms between (H^A, H_+^A) and (H^B, H_+^B) as ordered groups ([18]).

In this paper, we will study actions of the circle group $\mathbb{R}/\mathbb{Z}$ on the Cuntz–Krieger algebras from the view points of continuous orbit equivalence of one-sided topological Markov shifts and flow equivalence of two-sided topological Markov shifts. Let us denote by $\text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$ the set of actions of $\mathbb{R}/\mathbb{Z}$ on $\mathcal{O}_A$ trivially acting on the maximal abelian C^* -subalgebra $\mathcal{D}_A$. For $f \in C(X_A, \mathbb{Z})$, define the one-parameter unitary group $U_t(f)$, $t \in \mathbb{R}/\mathbb{Z}$ in $\mathcal{D}_A$ by

$$U_t(f) = \exp(2\pi\sqrt{-1}tf), \quad (1.6)$$

and an automorphism $\rho_t^{A,f}$ on $\mathcal{O}_A$ for each $t \in \mathbb{R}/\mathbb{Z}$ by

$$\rho_t^{A,f}(S_i) = U_t(f)S_i, \quad i = 1, \dots, N. \quad (1.7)$$

We know that $\rho_t^{A,f}$ belongs to $\text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$. For $\alpha \in \text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$, a continuous map $t \in \mathbb{R}/\mathbb{Z} \rightarrow u_t \in \mathcal{O}_A$ is called a unitary one-cocycle relative to α if $u_t, t \in \mathbb{R}/\mathbb{Z}$ are unitaries and satisfy $u_{t+s} = u_t \alpha_t(u_s)$, $t, s \in \mathbb{R}/\mathbb{Z}$. For $\alpha^1, \alpha^2 \in \text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$, we write $\alpha^1 \sim \alpha^2$ if there exists a unitary one-cocycle $u_t \in \mathcal{O}_A$ relative to α^2 such that $\alpha_t^1 = \text{Ad}_{u_t} \circ \alpha_t^2$ for all $t \in \mathbb{R}/\mathbb{Z}$. In Section 2, we show that the set $\text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$ has a natural structure of ordered group. We will then see that the map $f \in C(X_A, \mathbb{Z}) \rightarrow \rho^{A,f} \in \text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$ gives rise to an isomorphism of groups and induces an isomorphism from H^A onto $\text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)/\sim$ as ordered groups (Proposition 2.4). The action $\rho^{A,1}$ for the constant function 1 is the gauge action ρ^A which is an order unit of $\text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$. In Section 3, we will see that the map $\Psi_h : C(X_B, \mathbb{Z}) \rightarrow C(X_A, \mathbb{Z})$ defined in (1.5) controls actions of $\mathbb{R}/\mathbb{Z}$ on the Cuntz–Krieger algebras trivially acting on its maximal abelian C^* -subalgebras as in the following theorem.

Theorem 1.1 (Theorem 3.2). *Suppose that the one-sided topological Markov shifts (X_A, σ_A) and (X_B, σ_B) are continuously orbit equivalent via a homeomorphism $h : X_A \rightarrow X_B$. Then there exists an isomorphism $\Phi : \mathcal{O}_A \rightarrow \mathcal{O}_B$ satisfying the following conditions*

- (i) $\Phi(\mathcal{D}_A) = \mathcal{D}_B$,
- (ii) $\Phi \circ \rho_t^{A,f} = \rho_t^{B, \Psi_{h^{-1}}(f)} \circ \Phi$ for $f \in C(X_A, \mathbb{Z})$, $t \in \mathbb{R}/\mathbb{Z}$,
- (iii) $\Psi_{h^{-1}} : C(X_A, \mathbb{Z}) \rightarrow C(X_B, \mathbb{Z})$ induces an isomorphism from H^A to H^B as ordered abelian groups.

If one may take $l_1(x) = k_1(x) + 1, x \in X_A$ in (1.3) and $l_2(y) = k_2(y) + 1, y \in X_B$ in (1.4), then (X_A, σ_A) and (X_B, σ_B) are said to be *eventually one-sided conjugate*. By using above theorem, we then obtain the following result.

Theorem 1.2 (Corollary 3.5). *There exists an isomorphism $\Phi : \mathcal{O}_A \rightarrow \mathcal{O}_B$ such that*

$$\Phi(\mathcal{D}_A) = \mathcal{D}_B \quad \text{and} \quad \Phi \circ \rho_t^A = \rho_t^B \circ \Phi, \quad t \in \mathbb{R}/\mathbb{Z}$$

if and only if (X_A, σ_A) and (X_B, σ_B) are eventually one-sided conjugate.

For flow equivalence of two-sided topological Markov shifts, we have the following theorem by using Parry–Sullivan’s theorem [22] which says that the flow equivalence relation is generated by topological conjugacies of two-sided topological Markov shifts and expansions of the underlying directed graphs.

Theorem 1.3 (Theorem 5.7). *Suppose that the two-sided topological Markov shifts $(\bar{X}_A, \bar{\sigma}_A)$ and $(\bar{X}_B, \bar{\sigma}_B)$ are flow equivalent. Then there exist an isomorphism $\Phi : \mathcal{O}_A \otimes \mathcal{K} \rightarrow \mathcal{O}_B \otimes \mathcal{K}$, a homomorphism $\varphi : C(X_A, \mathbb{Z}) \rightarrow C(X_B, \mathbb{Z})$ of ordered abelian groups and a unitary one-cocycle $u_t^f \in \mathcal{U}(M(\mathcal{O}_A \otimes \mathcal{K}))$ relative to $\rho^{A,f} \otimes \text{id}$ for each function $f \in C(X_A, \mathbb{Z})$ satisfying the following conditions*

- (i) $\Phi(\mathcal{D}_A \otimes \mathcal{C}) = \mathcal{D}_B \otimes \mathcal{C}$,
- (ii) $\Phi \circ \text{Ad}(u_t^f) \circ (\rho_t^{A,f} \otimes \text{id}) = (\rho_t^{B, \varphi(f)} \otimes \text{id}) \circ \Phi$ for $f \in C(X_A, \mathbb{Z})$, $t \in \mathbb{R}/\mathbb{Z}$,
- (iii) $\varphi : C(X_A, \mathbb{Z}) \rightarrow C(X_B, \mathbb{Z})$ induces an isomorphism from H^A to H^B as ordered abelian groups.

In this paper, we denote by $\mathbb{N}$ the set of positive integers and by $\mathbb{Z}_+$ the set of nonnegative integers, respectively. For a topological Markov shift (X_A, σ_A) , a word $\mu = (\mu_1, \dots, \mu_k)$ for $\mu_i \in \{1, \dots, N\}$ is said to be admissible for X_A if there exists $(x_n)_{n \in \mathbb{N}} \in X_A$ such that $(\mu_1, \dots, \mu_k) = (x_1, \dots, x_k)$. The length of μ is k , which is denoted by $|\mu|$. We denote by $B_k(X_A)$ the set of all admissible words of length k . The cylinder set $\{(x_n)_{n \in \mathbb{N}} \in X_A \mid x_1 = \mu_1, \dots, x_k = \mu_k\}$ for $\mu = (\mu_1, \dots, \mu_k) \in B_k(X_A)$ is denoted by U_μ .

This paper is a revised version of the paper entitled *Continuous orbit equivalence, flow equivalence of Markov shifts and torus actions on Cuntz–Krieger algebras*, arXiv:1501.06965.

2 Circle actions

We fix an $N \times N$ irreducible matrix $A = [A(i, j)]_{i,j=1}^N$ with entries in $\{0, 1\}$. Let $S_1, \dots, S_N$ be the canonical generating partial isometries of the Cuntz-Krieger algebra $\mathcal{O}_A$ satisfying the relations (1.2). For a word $\mu = (\mu_1, \dots, \mu_k) \in B_k(X_A)$, denote by S_μ the partial isometry $S_{\mu_1} \cdots S_{\mu_k}$. Let us denote by $\text{Act}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$ the set of all continuous actions of the circle group $\mathbb{R}/\mathbb{Z}$ as automorphisms of $\mathcal{O}_A$. We set

$$\text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A) = \{\alpha \in \text{Act}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A) \mid \alpha_t(a) = a \text{ for all } t \in \mathbb{R}/\mathbb{Z}, a \in \mathcal{D}_A\}. \quad (2.1)$$

We denote by $\text{Rep}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$ (resp. $\text{Rep}(\mathbb{R}/\mathbb{Z}, \mathcal{D}_A)$) the set of unitary representations of $\mathbb{R}/\mathbb{Z}$ into the unitary group of $\mathcal{O}_A$ (resp. of $\mathcal{D}_A$). We are assuming that the matrix A is irreducible and not any permutation matrix, so that the subalgebra $\mathcal{D}_A$ is maximal abelian in $\mathcal{O}_A$ ([8]). Hence if $\alpha \in \text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$ is of the form $\alpha_t = \text{Ad}(u_t), t \in \mathbb{R}/\mathbb{Z}$ for some $u \in \text{Rep}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$, then $u_t \in \mathcal{D}_A$ for all $t \in \mathbb{R}/\mathbb{Z}$ and hence $u \in \text{Rep}(\mathbb{R}/\mathbb{Z}, \mathcal{D}_A)$.

Recall that for a function $f \in C(X_A, \mathbb{Z})$ and $t \in \mathbb{R}/\mathbb{Z}$, an automorphism $\rho_t^{A,f} \in \text{Aut}(\mathcal{O}_A)$ is defined by $\rho_t^{A,f}(S_i) = U_t(f)S_i, i = 1, \dots, N, t \in \mathbb{R}/\mathbb{Z}$ for the unitary $U_t(f) = \exp(2\pi\sqrt{-1}tf) \in \mathcal{D}_A$ as in (1.6) and (1.7). It is easy to see that the automorphisms $\rho_t^{A,f}, t \in \mathbb{R}/\mathbb{Z}$ yield an action of $\mathbb{R}/\mathbb{Z}$ on $\mathcal{O}_A$ such that $\rho_t^{A,f}(a) = a$ for all $a \in \mathcal{D}_A$ so that $\rho^{A,f} \in \text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$.

Lemma 2.1. *An action $\alpha \in \text{Act}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$ belongs to $\text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$ if and only if there exists a continuous function $f \in C(X_A, \mathbb{Z})$ such that $\alpha_t = \rho_t^{A,f}$ for all $t \in \mathbb{R}/\mathbb{Z}$.*

Proof. It suffices to show the only if part. Suppose that $\alpha \in \text{Act}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$ belongs to $\text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$. By [12, Lemma 4.6, Corollary 4.7], there exists a unitary $u_t \in \mathcal{U}(\mathcal{D}_A)$ for each $t \in \mathbb{R}/\mathbb{Z}$ such that $\alpha_t(S_i) = u_t S_i$ for $i = 1, \dots, N$ (cf. [7]). As $\alpha_s(u_t) = u_t$ for $s, t \in \mathbb{R}/\mathbb{Z}$, we see that for $i = 1, \dots, N$

$$u_t u_s S_i = u_t \alpha_s(S_i) = \alpha_s(u_t S_i) = \alpha_s(\alpha_t(S_i)) = \alpha_{s+t}(S_i) = u_{s+t} S_i$$

so that $u_t u_s = u_{s+t}$. The continuity of $u : \mathbb{R}/\mathbb{Z} \rightarrow \mathcal{U}(\mathcal{D}_A)$ follows from that of the action α . Hence there exists a unitary representation $u \in \text{Rep}(\mathbb{R}/\mathbb{Z}, \mathcal{D}_A)$ satisfying $\alpha_t(S_i) = u_t S_i$ for $i = 1, \dots, N$ and $t \in \mathbb{R}/\mathbb{Z}$. As in [16, Lemma 6.4], there exists a continuous function $f \in C(X_A, \mathbb{Z})$ such that $u_t = U_t(f)$ and hence $\alpha_t(S_i) = \rho_t^{A,f}(S_i)$ for $i = 1, \dots, N, t \in \mathbb{R}/\mathbb{Z}$. $\square$

For $\alpha, \beta \in \text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$, take $f, g \in C(X_A, \mathbb{Z})$ such that $\alpha_t(S_i) = U_t(f)S_i, \beta_s(S_i) = U_s(g)S_i, s, t \in \mathbb{R}/\mathbb{Z}, i = 1, \dots, N$. The commutativity $U_s(g)U_t(f) = U_t(f)U_s(g)$ implies $\alpha_t \circ \beta_s = \beta_s \circ \alpha_t$ for $s, t \in \mathbb{R}/\mathbb{Z}$. Hence the set $\text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$ has a structure of abelian group by the product $(\alpha \cdot \beta)_t = \alpha_t \circ \beta_t$ for $t \in \mathbb{R}/\mathbb{Z}$. As the set $C(X_A, \mathbb{Z})$ has a structure of abelian group by pointwise sums of functions, we get the following.

Proposition 2.2. *The correspondence $f \in C(X_A, \mathbb{Z}) \rightarrow \rho^{A,f} \in \text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$ yields an isomorphism of abelian groups.*

Proof. Since the identities

$$U_t(f_1 + f_2) = U_t(f_1)U_t(f_2), \quad U_t(-f) = U_t(f)^*, \quad U_t(0) = 1$$

hold for $f_1, f_2, f \in C(X_A, \mathbb{Z})$, we have

$$\rho_t^{A, f_1+f_2} = \rho_t^{A, f_1} \circ \rho_t^{A, f_2}, \quad \rho_t^{A, -f} = (\rho_t^{A, f})^{-1}, \quad \rho_t^{A, 0} = \text{id}.$$

Hence the correspondence $f \in C(X_A, \mathbb{Z}) \rightarrow \rho^{A, f} \in \text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$ yields a homomorphism of groups. Its surjectivity comes from the preceding lemma. It is clear to see that $\rho_t^{A, f} = \text{id}$ for all $t \in \mathbb{R}/\mathbb{Z}$ if and only if f is identically zero. $\square$

Under the identification between the algebras $C(X_A)$ and $\mathcal{D}_A$ through the correspondence between $\chi_{U_\mu} \in C(X_A)$ and $S_\mu S_\mu^* \in \mathcal{D}_A$ for $\mu \in B_k(X_A)$, we have $\sum_{i=1}^N S_i f S_i^* = f \circ \sigma_A$ for $f \in C(X_A)$ so that the equalities

$$S_i U_t(f) = \sum_{j=1}^N S_j U_t(f) S_j^* S_i = U_t(f \circ \sigma_A) S_i, \quad i = 1, \dots, N \quad (2.2)$$

hold. For a function $f \in C(X_A, \mathbb{Z})$, denote by $[f]$ the class of f in the ordered cohomology group H^A . We then have the following lemma.

Lemma 2.3. *For $f_1, f_2 \in C(X_A, \mathbb{Z})$, the following two conditions are equivalent:*

- (i) $[f_1] = [f_2]$ in H^A .
- (ii) $\rho_t^{A, f_2} = \text{Ad}(u_t) \circ \rho_t^{A, f_1}$ on $\mathcal{O}_A$ for some $u \in \text{Rep}(\mathbb{R}/\mathbb{Z}, \mathcal{D}_A)$.

Proof. (i) $\Rightarrow$ (ii): Since $[f_1] = [f_2]$ in H^A , there exists a continuous function $b \in C(X_A, \mathbb{Z})$ such that $f_2 = f_1 + b - b \circ \sigma_A$. Put $u_t = U_t(b) \in \mathcal{U}(\mathcal{D}_A)$ for $t \in \mathbb{R}/\mathbb{Z}$. By the identity (2.2), one sees that

$$u_t \rho_t^{f_1}(S_i) u_t^* = U_t(b) U_t(f_1) S_i U_t(b)^* = U_t(b) U_t(f_1) U_t(-b \circ \sigma_A) S_i = U_t(f_2) S_i$$

so that we have $\text{Ad}(u_t) \circ \rho_t^{A, f_1} = \rho_t^{A, f_2}$.

(ii) $\Rightarrow$ (i): By [16, Lemma 6.4], there exists a continuous function $b \in C(X_A, \mathbb{Z})$ such that $u_t = U_t(b)$. The equality $\rho_t^{A, f_2}(S_i) = \text{Ad}(u_t) \circ \rho_t^{A, f_1}(S_i)$ implies that

$$U_t(f_2) S_i = U_t(b + f_1 - b \circ \sigma_A) S_i, \quad i = 1, \dots, N$$

so that $f_2 = f_1 + b - b \circ \sigma_A$ and hence $[f_1] = [f_2]$. $\square$

For $\alpha^1, \alpha^2 \in \text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$, we write $\alpha^1 \sim \alpha^2$ if there exists a unitary one-cocycle $u_t \in \mathcal{O}_A, t \in \mathbb{R}/\mathbb{Z}$ relative to α^2 such that $\alpha_t^1 = \text{Ad} u_t \circ \alpha_t^2$ for $t \in \mathbb{R}/\mathbb{Z}$. In this case, we know that u_t belongs to $\mathcal{D}_A$ and satisfies $u_{t+s} = u_t u_s$ for $t, s \in \mathbb{R}/\mathbb{Z}$ because $\alpha_t^i(a) = a$ for all $a \in \mathcal{D}_A$ and $\mathcal{D}_A$ is maximal abelian in $\mathcal{O}_A$. It is easy to see that $\sim$ is an equivalence relation in $\text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$. We write $[\alpha]$ for the equivalence class of $\alpha \in \text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$. For $\alpha \in \text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$, we put

$$\delta(\alpha)_t = \frac{d}{dt} \left(\sum_{i=1}^N \alpha_t(S_i) S_i^* \right) (t).$$

Define $\alpha \geq 0$ if $\frac{1}{2\pi\sqrt{-1}} \delta(\alpha)_0$ is a positive operator in $\mathcal{O}_A$. We write $[\alpha] \geq 0$ if there exists $\alpha' \in \text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$ such that $\alpha' \geq 0$ and $\alpha \sim \alpha'$. As $\delta(\rho^{A, f})_0 = 2\pi\sqrt{-1}f$, we have

Proposition 2.4. *The isomorphism $f \in C(X_A, \mathbb{Z}) \rightarrow \rho^{A, f} \in \text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A)$ of groups induces an isomorphism $[f] \in H^A \rightarrow [\rho^{A, f}] \in \text{Act}_{\mathcal{D}_A}(\mathbb{R}/\mathbb{Z}, \mathcal{O}_A) / \sim$ of ordered abelian groups.*

3 Continuous orbit equivalence and circle actions

Let $h : X_A \rightarrow X_B$ be a homeomorphism which gives rise to a continuous orbit equivalence between (X_A, σ_A) and (X_B, σ_B) . Take $k_1, l_1 \in C(X_A, \mathbb{Z}_+)$ satisfying (1.3). For $f \in C(X_B, \mathbb{Z})$, define $\Psi_h(f) \in C(X_A, \mathbb{Z})$ by the formula (1.5). The map $\Psi_h : C(X_B, \mathbb{Z}) \rightarrow C(X_A, \mathbb{Z})$ does not depend on the choice of the functions k_1, l_1 as long as they are satisfying (1.3) ([18, Lemma 4.2]). Thus $\Psi_h : C(X_B, \mathbb{Z}) \rightarrow C(X_A, \mathbb{Z})$ gives rise to a homomorphism of abelian groups, which is actually an isomorphism ([18, Proposition 4.5]). As the equality $\Psi_h(f - f \circ \sigma_B) = f \circ h - f \circ h \circ \sigma_A$ holds, Ψ_h induces a homomorphism from H^B to H^A which yields an isomorphism of ordered groups ([17], [18]). For $f \in C(X_A, \mathbb{Z})$ and $n \in \mathbb{Z}_+$, let us denote by f^n the function $f^n(x) = \sum_{i=0}^{n-1} f(\sigma_A^i(x))$, $x \in X_A$. The following identity is useful in our further discussions.

Lemma 3.1. *For a word $\mu = (\mu_1, \dots, \mu_n) \in B_n(X_A)$, we have*

$$\rho_t^{A,f}(S_\mu) = U_t(f^n)S_\mu, \quad f \in C(X_A, \mathbb{Z}), t \in \mathbb{R}/\mathbb{Z}. \quad (3.1)$$

Proof. By the identity $S_i U_t(f) = U_t(f \circ \sigma_A) S_i$ as in (2.2), we have

$$\begin{aligned} \rho_t^{A,f}(S_{\mu_1 \dots \mu_n}) &= U_t(f)S_{\mu_1} \cdots U_t(f)S_{\mu_{n-2}}U_t(f)S_{\mu_{n-1}}U_t(f)S_{\mu_n} \\ &= U_t(f)S_{\mu_1} \cdots U_t(f)S_{\mu_{n-2}}U_t(f + f \circ \sigma_A)S_{\mu_{n-1}}S_{\mu_n} \\ &= U_t(f + f \circ \sigma_A + \cdots + f \circ \sigma_A^{n-1})S_{\mu_1} \cdots S_{\mu_{n-2}}S_{\mu_{n-1}}S_{\mu_n} \\ &= U_t(f^n)S_\mu. \end{aligned}$$

□

The following result is a generalization of [8, 2.17 Proposition].

Theorem 3.2. *If the one-sided topological Markov shifts (X_A, σ_A) and (X_B, σ_B) are continuously orbit equivalent via a homeomorphism $h : X_A \rightarrow X_B$, then there exists an isomorphism $\Phi : \mathcal{O}_A \rightarrow \mathcal{O}_B$ such that*

$$\Phi(\mathcal{D}_A) = \mathcal{D}_B \quad \text{and} \quad \Phi \circ \rho_t^{A, \Psi_h(f)} = \rho_t^{B,f} \circ \Phi \quad \text{for } f \in C(X_B, \mathbb{Z}), t \in \mathbb{R}/\mathbb{Z}.$$

Proof. The first part of the proof below follows essentially the proof of [14, Proposition 5.6.] (cf. [16, Proposition 6.3]). We need the several notations used in the first part to proceed to the second part, so that the first part overlaps [14, Proposition 5.6.] and [16, Proposition 6.3]. Let us denote by $\mathfrak{H}_A$ (resp. $\mathfrak{H}_B$) the Hilbert space with its complete orthonormal system $\{e_x^A \mid x \in X_A\}$ (resp. $\{e_y^B \mid y \in X_B\}$). Define the partial isometries $S_i^A, i = 1, \dots, N$ on $\mathfrak{H}_A$ by

$$S_i^A e_x^A = \begin{cases} e_{ix}^A & \text{if } ix \in X_A, \\ 0 & \text{otherwise,} \end{cases} \quad (3.2)$$

which satisfy the relations (1.2). For the $M \times M$ matrix $B = [B(i, j)]_{i,j=1}^M$, define the partial isometries $S_i^B, i = 1, \dots, M$ on $\mathfrak{H}_B$ satisfying the relations (1.2) for B in a similar way. The Cuntz–Krieger algebra $\mathcal{O}_A$ (resp. $\mathcal{O}_B$) is identified with the C^* -algebra $C^*(S_1^A, \dots, S_N^A)$ (resp. $C^*(S_1^B, \dots, S_M^B)$) on $\mathfrak{H}_A$ (resp. $C^*(S_1^B, \dots, S_M^B)$ on $\mathfrak{H}_B$) generated by $S_1^A, \dots, S_N^A$ (resp. $S_1^B, \dots, S_M^B$). For

the continuous function $k_1 : X_A \rightarrow \mathbb{Z}_+$ in (1.3), let $K_1 = \text{Max}\{k_1(x) \mid x \in X_A\}$. By adding $K_1 - k_1(x)$ to both $k_1(x)$ and $l_1(x)$, one may assume that $k_1(x) = K_1$ for all $x \in X_A$.

We will see that $\Phi = \text{Ad}(U_h)$ defined by the unitary $U_h e_x^A = e_{h(x)}^B, x \in X_A$ satisfies the desired properties. We fix $i \in \{1, \dots, N\}$ and set $X_B^{(i)} = \{y \in X_B \mid ih^{-1}(y) \in X_A\}$. For $y \in X_B^{(i)}$, put $z = ih^{-1}(y) \in X_A$. As $h(\sigma_A(z)) = y$ with (1.3), one has $h(z) \in \sigma_B^{-l_1(z)}(\sigma_B^{K_1}(y))$ and

$$h(z) = (\mu_1(z), \dots, \mu_{l_1(z)}(z), y_{K_1+1}, y_{K_1+2}, \dots) \quad (3.3)$$

for some $(\mu_1(z), \dots, \mu_{l_1(z)}(z)) \in B_{l_1(z)}(X_B)$. Put $L_1 = \text{Max}\{l_1(z) \mid z = ih^{-1}(y), y \in X_B^{(i)}\}$. The set $W^{(i)} = \{(\mu_1(z), \dots, \mu_{l_1(z)}(z)) \in B_{l_1(z)}(X_B) \mid z = ih^{-1}(y), y \in X_B^{(i)}\}$ of words is a finite subset of $\cup_{j=0}^{L_1} B_j(X_B)$. For $\nu = (\nu_1, \dots, \nu_j) \in W^{(i)}$, put the clopen set $E_\nu^{(i)}$ in $X_B^{(i)}$

$$E_\nu^{(i)} = \{y \in X_B^{(i)} \mid \mu_1(z) = \nu_1, \dots, \mu_{l_1(z)}(z) = \nu_j, z = ih^{-1}(y)\}$$

so that $X_B^{(i)} = \cup_{\nu \in W^{(i)}} E_\nu^{(i)}$. Let $Q_\nu^{(i)}$ be the characteristic function $\chi_{E_\nu^{(i)}}$ on X_B for the clopen sets $E_\nu^{(i)}$. For $y \in X_B^{(i)}$ and $\nu \in W^{(i)}$, we have $y \in E_\nu^{(i)}$ if and only if $Q_\nu^{(i)} e_y^B = e_y^B$. By (3.3), the equality

$$e_{h(ih^{-1}(y))}^B = \sum_{\nu \in W^{(i)}} S_\nu^B \sum_{\xi \in B_{K_1}(X_B)} S_\xi^{B*} Q_\nu^{(i)} e_y^B \quad \text{for } y \in X_B^{(i)}$$

holds. As $U_h S_i^A U_h^* e_y^B = e_{h(ih^{-1}(y))}^B$ if $y \in X_B^{(i)}$, and 0 otherwise, we have

$$U_h S_i^A U_h^* e_y^B = \sum_{\nu \in W^{(i)}} \sum_{\xi \in B_{K_1}(X_B)} S_\nu^B S_\xi^{B*} Q_\nu^{(i)} e_y^B \quad \text{for } y \in X_B^{(i)}$$

so that

$$U_h S_i^A U_h^* = \sum_{\nu \in W^{(i)}} \sum_{\xi \in B_{K_1}(X_B)} S_\nu^B S_\xi^{B*} Q_\nu^{(i)}.$$

As $Q_\nu^{(i)} \in \mathcal{D}_B$, we have $\text{Ad}(U_h)(S_i^A) \in \mathcal{O}_B$ so that $\text{Ad}(U_h)(\mathcal{O}_A) \subset \mathcal{O}_B$. Since $U_h^* = U_{h^{-1}}$, we symmetrically have $\text{Ad}(U_h^*)(\mathcal{O}_B) \subset \mathcal{O}_A$ so that $\text{Ad}(U_h)(\mathcal{O}_A) = \mathcal{O}_B$. It is direct to see that $\text{Ad}(U_h)(f) = f \circ h^{-1}$ for $f \in \mathcal{D}_A$ from the definition $U_h e_x^A = e_{h(x)}^B, x \in X_A$ so that we have $\text{Ad}(U_h)(\mathcal{D}_A) = \mathcal{D}_B$.

We will next show that $\text{Ad}(U_h) \circ \rho_t^{A, \Psi_h(f)} = \rho_t^{B, f} \circ \text{Ad}(U_h)$ for $f \in C(X_B, \mathbb{Z}), t \in \mathbb{R}/\mathbb{Z}$. It follows that

$$(\rho_t^{B, f} \circ \text{Ad}(U_h))(S_i^A) = \sum_{\nu \in W^{(i)}} \sum_{\xi \in B_{K_1}(X_B)} \rho_t^{B, f}(S_\nu^B S_\xi^{B*}) Q_\nu^{(i)}. \quad (3.4)$$

Since $Q_\nu^{(i)} e_y^B \neq 0$ if and only if $Q_\nu^{(i)} e_y^B = e_y^B$ and $\nu_1 = \mu_1(z), \dots, \nu_j = \mu_{l_1(z)}(z)$. For $y \in E_\nu^{(i)}$ with $(y_1, \dots, y_{K_1}) = (\xi_1, \dots, \xi_{K_1})$, we have $S_\nu^B S_\xi^{B*} Q_\nu^{(i)} e_y^B = e_{h(z)}^B = e_{h(ih^{-1}(y))}^B$ and $\nu \sigma_B^{K_1}(y) = h(z)$. As

$$\rho_t^{B, f}(S_\xi^{B*}) e_y^B = \exp\{-2\pi\sqrt{-1}t f^{K_1}(y)\} S_\xi^{B*} e_y^B = \exp\{-2\pi\sqrt{-1}t f^{K_1}(h(\sigma_A(z)))\} e_{\sigma_B^{K_1}(y)}^B$$

and

$$\rho_t^{B,f}(S_\nu^B)e_{\sigma_B^{K_1}(y)}^B = U_t(f^{l_1(z)})e_{\nu\sigma_B^{K_1}(y)}^B = \exp\{2\pi\sqrt{-1}tf^{l_1(z)}(h(z))\}e_{h(z)}^B,$$

we have

$$\begin{aligned} \rho_t^{B,f}(S_\nu^B S_\xi^{B*} Q_\nu^{(i)})e_y^B &= \rho_t^{B,f}(S_\nu^B)\rho_t^{B,f}(S_\xi^{B*})e_y^B \\ &= \rho_t^{B,f}(S_\nu^B)\exp\{-2\pi\sqrt{-1}tf^{K_1}(h(\sigma_A(z)))\}e_{\sigma_B^{K_1}(y)}^B \\ &= \exp\{-2\pi\sqrt{-1}tf^{K_1}(h(\sigma_A(z)))\}\rho_t^{B,f}(S_\nu^B)e_{\sigma_B^{K_1}(y)}^B \\ &= \exp\{-2\pi\sqrt{-1}tf^{K_1}(h(\sigma_A(z)))\}\exp\{2\pi\sqrt{-1}tf^{l_1(z)}(h(z))\}e_{h(z)}^B \\ &= \exp\{2\pi\sqrt{-1}t(\Psi_h(f)(z))\}e_{h(z)}^B \\ &= U_h\exp\{2\pi\sqrt{-1}t\Psi_h(f)\}S_i^A e_{h^{-1}(y)}^A \\ &= U_h\rho_t^{A,\Psi_h(f)}(S_i^A)U_h^*e_y^B. \end{aligned}$$

By (3.4), we get

$$(\rho_t^{B,f} \circ \text{Ad}(U_h))(S_i^A)e_y^B = U_h\rho_t^{A,\Psi_h(f)}(S_i^A)U_h^*e_y^B$$

so that

$$\rho_t^{B,f} \circ \text{Ad}(U_h) = \text{Ad}(U_h) \circ \rho_t^{A,\Psi_h(f)} \quad \text{for } t \in \mathbb{R}/\mathbb{Z}.$$

By setting $\Phi = \text{Ad}(U_h) : \mathcal{O}_A \rightarrow \mathcal{O}_B$, we have the desired isomorphism. $\square$

In the proof of [14, Theorem 5.7] we see that if there exists an isomorphism $\Phi : \mathcal{O}_A \rightarrow \mathcal{O}_B$ such that $\Phi(\mathcal{D}_A) = \mathcal{D}_B$, we can take a homeomorphism $h : X_A \rightarrow X_B$ which gives rise to a continuous orbit equivalence between (X_A, σ_A) and (X_B, σ_B) and satisfies $\Phi(f) = f \circ h^{-1}$ for $f \in \mathcal{D}_A$. We apply Theorem 3.2 to obtain the following theorem.

Theorem 3.3. *Let $f \in C(X_B, \mathbb{Z}), g \in C(X_A, \mathbb{Z})$.*

(i) *There exists an isomorphism $\Phi : \mathcal{O}_A \rightarrow \mathcal{O}_B$ satisfying*

$$\Phi(\mathcal{D}_A) = \mathcal{D}_B \quad \text{and} \quad \Phi \circ \rho_t^{A,g} = \rho_t^{B,f} \circ \Phi, \quad t \in \mathbb{R}/\mathbb{Z} \quad (3.5)$$

if and only if there exists a homeomorphism $h : X_A \rightarrow X_B$ which gives rise to a continuous orbit equivalence between (X_A, σ_A) and (X_B, σ_B) such that $\Psi_h(f) = g$.

(ii) *There exist an isomorphism $\Phi : \mathcal{O}_A \rightarrow \mathcal{O}_B$ and a unitary one-cocycle $v_t, t \in \mathbb{R}/\mathbb{Z}$ in $\mathcal{O}_A$ relative to $\rho^{A,g}$ satisfying*

$$\Phi(\mathcal{D}_A) = \mathcal{D}_B \quad \text{and} \quad \Phi \circ \text{Ad}(v_t) \circ \rho_t^{A,g} = \rho_t^{B,f} \circ \Phi, \quad t \in \mathbb{R}/\mathbb{Z} \quad (3.6)$$

if and only if there exists a homeomorphism $h : X_A \rightarrow X_B$ which gives rise to a continuous orbit equivalence between (X_A, σ_A) and (X_B, σ_B) such that $[\Psi_h(f)] = [g]$ in H^A .

Proof. (i) The if part follows from Theorem 3.2. We will show the only if part. Represent the algebras $\mathcal{O}_A$ on $\mathfrak{H}_A$ and $\mathcal{O}_B$ on $\mathfrak{H}_B$ as in the proof of Theorem 3.2. Suppose that there exists an isomorphism $\Phi : \mathcal{O}_A \rightarrow \mathcal{O}_B$ satisfying (3.5). Take a homeomorphism

$h : X_A \rightarrow X_B$ which gives rise to a continuous orbit equivalence between (X_A, σ_A) and (X_B, σ_B) and satisfies $\Phi(f) = f \circ h^{-1}$ for $f \in \mathcal{D}_A$. Let $\Phi_h : \mathcal{O}_A \rightarrow \mathcal{O}_B$ be the isomorphism $\text{Ad}(U_h)$ defined in the proof of Theorem 3.2 so that

$$\Phi_h(\mathcal{D}_A) = \mathcal{D}_B \quad \text{and} \quad \Phi_h \circ \rho_t^{A, \Psi_h(f)} = \rho_t^{B, f} \circ \Phi_h, \quad t \in \mathbb{R}/\mathbb{Z}. \quad (3.7)$$

By the construction of Φ_h , we know that $\Phi_h(f) = f \circ h^{-1}$ for $f \in \mathcal{D}_A$. Hence the automorphism $\alpha = \Phi_h^{-1} \circ \Phi$ on $\mathcal{O}_A$ satisfies $\alpha|_{\mathcal{D}_A} = \text{id}$. By [14, Theorem 6.5 (1)], there exists a unitary one-cocycle V_α in $\mathcal{D}_A$ relative to α such that $\alpha(S_\mu) = V_\alpha(k)S_\mu$ for $\mu \in B_k(X_A)$. Hence $\alpha(S_i) = V_\alpha(1)S_i$, $V_\alpha(1) \in \mathcal{D}_A$ so that $\Phi(S_i) = \Phi_h(V_\alpha(1)S_i)$ for $i = 1, \dots, N$. By (3.5) and (3.7), we see

$$\Phi \circ \rho_t^{A, g}(S_i) = (\Phi_h \circ \rho_t^{A, \Psi_h(f)} \circ \Phi_h^{-1}) \circ \Phi(S_i), \quad t \in \mathbb{R}/\mathbb{Z}, i = 1, \dots, N.$$

The left hand side above equals

$$\Phi(\rho_t^{A, g}(S_i)) = \Phi(U_t(g)S_i) = \Phi(U_t(g))\Phi(S_i).$$

The right hand side above equals

$$\begin{aligned} & (\Phi_h \circ \rho_t^{A, \Psi_h(f)} \circ \Phi_h^{-1}) \circ \Phi(S_i) \\ &= \Phi_h(V_\alpha(1)\rho_t^{A, \Psi_h(f)}(S_i)) = \Phi_h(U_t(\Psi_h(f)))\Phi_h(V_\alpha(1)S_i) = \Phi_h(U_t(\Psi_h(f)))\Phi(S_i). \end{aligned}$$

Hence we have $\Phi(U_t(g))\Phi(S_i) = \Phi_h(U_t(\Psi_h(f)))\Phi(S_i)$. As both $U_t(g), U_t(\Psi_h(f))$ belong to $\mathcal{D}_A$ and $\Phi = \Phi_h$ on $\mathcal{D}_A$, we obtain that $U_t(g) = U_t(\Psi_h(f))$ and hence $g = \Psi_h(f)$.

(ii) We first show the if part. Suppose that $[\Psi_h(f)] = [g]$. We may take $b \in C(X_A, \mathbb{Z})$ such that $\Psi_h(f) = g + b - b \circ \sigma_A$. By (i), there exists an isomorphism $\Phi : \mathcal{O}_A \rightarrow \mathcal{O}_B$ such that $\Phi(\mathcal{D}_A) = \mathcal{D}_B$ and $\Phi \circ \rho_t^{A, g+b-b \circ \sigma_A} = \rho_t^{B, f} \circ \Phi$. Since the unitary $v_t = U_t(b)$ satisfies $\text{Ad}(v_t) \circ \rho_t^{A, g} = \rho_t^{A, g+b-b \circ \sigma_A}$, the equality (3.6) holds.

For the proof of the only if part, assume that there exist an isomorphism $\Phi : \mathcal{O}_A \rightarrow \mathcal{O}_B$ and a unitary one-cocycle $v_t, t \in \mathbb{R}/\mathbb{Z}$ in $\mathcal{O}_A$ relative to $\rho^{A, g}$ satisfying the equality (3.6). As $\rho_t^{A, g}|_{\mathcal{D}_A} = \text{id}$ and $\rho_t^{B, f}|_{\mathcal{D}_B} = \text{id}$, the equality (3.6) implies that $\Phi(v_t)$ commutes with elements of $\Phi(\mathcal{D}_A) = \mathcal{D}_B$. Since $\mathcal{D}_B$ is maximal abelian in $\mathcal{O}_B$, the unitaries $\Phi(v_t), t \in \mathbb{R}/\mathbb{Z}$ belong to $\mathcal{D}_B$ and hence $v_t, t \in \mathbb{R}/\mathbb{Z}$ belong to $\mathcal{D}_A$ which satisfy $\rho_s^{A, g}(v_t) = v_t$ for $s, t \in \mathbb{R}/\mathbb{Z}$. The map $t \in \mathbb{R}/\mathbb{Z} \rightarrow v_t \in \mathcal{D}_A$ then yields a unitary representation of $\mathbb{R}/\mathbb{Z}$. Take $b \in C(X_A, \mathbb{Z})$ such that $v_t = U_t(b)$ and hence the equality $\Phi \circ \rho_t^{A, g+b-b \circ \sigma_A} = \rho_t^{B, f} \circ \Phi$ follows. By (i), there exists a homeomorphism $h : X_A \rightarrow X_B$ which gives rise to a continuous orbit equivalence between (X_A, σ_A) and (X_B, σ_B) such that $\Phi_h(f) = g + b - b \circ \sigma_A$ so that $[\Psi_h(f)] = [g]$. $\square$

Under the assumption of Theorem 3.2, let us define the functions $c_1 \in C(X_A, \mathbb{Z})$ and $c_2 \in C(X_B, \mathbb{Z})$ by $c_1(x) = l_1(x) - k_1(x), x \in X_A$ and $c_2(y) = l_2(y) - k_2(y), y \in X_B$, respectively. They are called the cocycle functions which play a key rôle in [16] and [18]. They do not depend on the choices of the functions k_1, l_1 and k_2, l_2 as long as they are satisfying (1.3) and (1.4), respectively ([18, Lemma 4.1]). By the formula (1.5), we know $\Psi_h(1_B) = c_1$ and similarly $\Psi_{h^{-1}}(1_A) = c_2$, where $1_A, 1_B$ are the constant functions on X_A, X_B taking its values 1, respectively. The actions $\rho_t^{A, 1_A}$ for the constant function 1_A and $\rho_t^{B, 1_B}$ for the constant function 1_B are the gauge actions ρ_t^A on $\mathcal{O}_A$ and ρ_t^B on $\mathcal{O}_B$, respectively. Theorem 3.3 (i) implies the following corollary.

Corollary 3.4. *The one-sided topological Markov shifts (X_A, σ_A) and (X_B, σ_B) are continuously orbit equivalent if and only if there exist an isomorphism $\Phi : \mathcal{O}_A \rightarrow \mathcal{O}_B$ and continuous functions $c_1 : X_A \rightarrow \mathbb{Z}$, $c_2 : X_B \rightarrow \mathbb{Z}$, such that*

$$\Phi(\mathcal{D}_A) = \mathcal{D}_B \quad \text{and} \quad \Phi \circ \rho_t^A = \rho_t^{B, c_2} \circ \Phi, \quad \Phi^{-1} \circ \rho_t^B = \rho_t^{A, c_1} \circ \Phi \quad \text{for } t \in \mathbb{R}/\mathbb{Z}.$$

In the case $f = 1_B, g = 1_A$ in Theorem 3.3, the first assertion (i) describes the commutativity of the gauge actions. We say (X_A, σ_A) and (X_B, σ_B) to be *eventually one-sided conjugate* if there exist a homeomorphism $h : X_A \rightarrow X_B$ and continuous functions $k_1 : X_A \rightarrow \mathbb{Z}$ and $k_2 : X_B \rightarrow \mathbb{Z}$ such that

$$\sigma_B^{k_1(x)}(h(\sigma_A(x))) = \sigma_B^{k_1(x)+1}(h(x)) \quad \text{for } x \in X_A, \quad (3.8)$$

$$\sigma_A^{k_2(y)}(h^{-1}(\sigma_B(y))) = \sigma_A^{k_2(y)+1}(h^{-1}(y)) \quad \text{for } y \in X_B. \quad (3.9)$$

If the conditions (3.8) and (3.9) are fulfilled, one may take the functions k_1, k_2 to be the constant functions taking its values $K = \max\{k_1(x), k_2(y) \mid x \in X_A, y \in X_B\}$. As $\Psi_{h^{-1}} = (\Psi_h)^{-1}$, the condition $\Psi_h(1_B) = 1_A$ is equivalent to the condition $\Psi_{h^{-1}}(1_A) = 1_B$. The latter and hence the former conditions are equivalent to the conditions $l_1(x) = k_1(x) + 1$ and $l_2(y) = k_2(y) + 1$ by the definition of $\Psi_h, \Psi_{h^{-1}}$. This means that the eventual one-sided conjugacy is equivalent to the condition $\Psi_h(1_B) = 1_A$. As a more general equivalence relation, a notion of strongly continuous orbit equivalence between one-sided topological Markov shifts has been introduced in [16] which is stronger than continuous orbit equivalence but weaker than eventual one-sided conjugacy. Two one-sided topological Markov shifts (X_A, σ_A) and (X_B, σ_B) are said to be strongly continuous orbit equivalent if there exists a homeomorphism $h : X_A \rightarrow X_B$ giving rise to a continuous orbit equivalence such that $[\Psi_h(1_B)] = [1_A]$ in H^A , which is equivalent to the condition $[\Psi_{h^{-1}}(1_A)] = [1_B]$ in H^B ([16, Lemma 4.3]). We then have the following corollary of Theorem 3.3.

Corollary 3.5 (cf. [8, 2.17. Proposition], [16, Theorem 6.7]). *Let ρ^A (resp. ρ^B) be the gauge action on $\mathcal{O}_A$ (resp. $\mathcal{O}_B$).*

- (i) *There exists an isomorphism $\Phi : \mathcal{O}_A \rightarrow \mathcal{O}_B$ such that*

$$\Phi(\mathcal{D}_A) = \mathcal{D}_B \quad \text{and} \quad \Phi \circ \rho_t^A = \rho_t^B \circ \Phi, \quad t \in \mathbb{R}/\mathbb{Z}$$

if and only if (X_A, σ_A) and (X_B, σ_B) are eventually one-sided conjugate.

- (ii) *There exist an isomorphism $\Phi : \mathcal{O}_A \rightarrow \mathcal{O}_B$ and a unitary representation v of $\mathbb{R}/\mathbb{Z}$ to $\mathcal{D}_B$ such that*

$$\Phi(\mathcal{D}_A) = \mathcal{D}_B \quad \text{and} \quad \Phi \circ \rho_t^A = \text{Ad}(v_t) \circ \rho_t^B \circ \Phi, \quad t \in \mathbb{R}/\mathbb{Z}$$

if and only if (X_A, σ_A) and (X_B, σ_B) are strongly continuous orbit equivalent.

Remark 3.6. Cuntz and Krieger had proved that if (X_A, σ_A) and (X_B, σ_B) are one-sided conjugate, then there exists an isomorphism $\Phi : \mathcal{O}_A \rightarrow \mathcal{O}_B$ such that $\Phi(\mathcal{D}_A) = \mathcal{D}_B$ and $\Phi \circ \rho_t^A = \rho_t^B \circ \Phi, t \in \mathbb{R}/\mathbb{Z}$ ([8, 2.17. Proposition]). The converse implication is seen in [29] as an open question. By the above corollary, the open question may be rephrased such that whether or not the eventual one-sided conjugacy implies one-sided conjugacy. This seems to be open.

We denote by $C(X_A, \mathbb{R})$ the set of real valued continuous functions on X_A . Let $\gamma_r^{A,f} \in \text{Aut}(\mathcal{O}_A)$, $r \in \mathbb{R}$ be the one-parameter automorphism for $f \in C(X_A, \mathbb{R})$ on $\mathcal{O}_A$ defined by

$$\gamma_r^{A,f}(S_i) = \exp(\sqrt{-1}rf)S_i, \quad i = 1, \dots, N. \quad (3.10)$$

For a homeomorphism $h : X_A \rightarrow X_B$ which gives rise to a continuous orbit equivalence between (X_A, σ_A) and (X_B, σ_B) , we define a linear map $\Psi_h : C(X_B, \mathbb{R}) \rightarrow C(X_A, \mathbb{R})$ by the same formula as in (1.5). We may show the following proposition in a similar way to Theorem 3.2.

Proposition 3.7. *If the one-sided topological Markov shifts (X_A, σ_A) and (X_B, σ_B) are continuously orbit equivalent via a homeomorphism $h : X_A \rightarrow X_B$, then there exists an isomorphism $\Phi : \mathcal{O}_A \rightarrow \mathcal{O}_B$ such that*

$$\Phi(\mathcal{D}_A) = \mathcal{D}_B \quad \text{and} \quad \Phi \circ \gamma_r^{A, \Psi_h(f)} = \gamma_r^{B, f} \circ \Phi \quad \text{for } f \in C(X_B, \mathbb{R}), \quad r \in \mathbb{R}.$$

4 Strong shift equivalence and circle actions

It is well-known that a topological Markov shift defined by a square matrix A with entries in $\{0, 1\}$ is naturally identified with a shift of finite type of the edge shift defined by the underlying directed graph. In this section, we consider edge shifts and hence square matrices with entries in nonnegative integers (cf. [10], [11], [30], etc.). Such a matrix is simply called a nonnegative square matrix. Our hypothesis that the shift space X_A is homeomorphic to a Cantor discontinuum forces the nonnegative square matrix to be irreducible and not any permutation matrix. For a nonnegative square matrix A , denote by E_A the edge set of the underlying graph G_A . The two-sided shift space $\bar{X}_A$ is identified with the two-sided sequences of concatenated edges of E_A . Two square matrices A, B are said to be elementary equivalent if there exist nonnegative rectangular matrices C, D such that $A = CD$ and $B = DC$ ([30], cf. [11]). The graphs G_A and G_B become both bipartite graphs. We may then take certain bijections $\varphi_{A,CD}$ from E_A to a subset of $E_C \times E_D$ and $\varphi_{B,DC}$ from E_B to a subset of $E_D \times E_C$. We fix such bijections. Through the bijections, we may identify an edge a of E_A with a pair cd of edges $c \in E_C$ and $d \in E_D$, and similarly an edge b of E_B with a pair dc of edges $d \in E_D$ and $c \in E_C$. An equivalence relation generated by elementary equivalences is called the strong shift equivalence. In [30], R. Williams has proved that two-sided topological Markov shifts $(\bar{X}_A, \bar{\sigma}_A)$ and $(\bar{X}_B, \bar{\sigma}_B)$ for irreducible matrices A and B are topologically conjugate if and only if the matrices A and B are strong shift equivalent.

For a C^* -algebra $\mathcal{A}$ without unit, let $M(\mathcal{A})$ stand for its multiplier C^* -algebra defined by

$$M(\mathcal{A}) = \{a \in \mathcal{A}^{**} \mid a\mathcal{A} \subset \mathcal{A}, \mathcal{A}a \subset \mathcal{A}\}.$$

An action α of $\mathbb{R}/\mathbb{Z}$ on $\mathcal{A}$ always extends to $M(\mathcal{A})$ and we write the extended action still α . For an action α of $\mathbb{R}/\mathbb{Z}$ on $\mathcal{A}$, a unitary one-cocycle $u_t, t \in \mathbb{R}/\mathbb{Z}$ relative to α is a continuous map $t \in \mathbb{R}/\mathbb{Z} \rightarrow u_t \in \mathcal{U}(M(\mathcal{A}))$ to the unitary group $\mathcal{U}(M(\mathcal{A}))$ satisfying $u_{t+s} = u_s \alpha_s(u_t)$, $s, t \in \mathbb{R}/\mathbb{Z}$. It is known that if two matrices A and B are elementary equivalent, there exists an isomorphism $\Phi : \mathcal{O}_A \otimes \mathcal{K} \rightarrow \mathcal{O}_B \otimes \mathcal{K}$ and a unitary one-cocycle $u_t \in \mathcal{U}(M(\mathcal{O}_A \otimes \mathcal{K}))$ relative to the gauge action $\rho^A \otimes \text{id}$ such that

$$\Phi(\mathcal{D}_A \otimes \mathcal{C}) = \mathcal{D}_B \otimes \mathcal{C} \quad \text{and} \quad \Phi \circ \text{Ad}(u_t) \circ (\rho_t^A \otimes \text{id}) = (\rho_t^B \otimes \text{id}) \circ \Phi \quad \text{for } t \in \mathbb{R}/\mathbb{Z}$$

(cf. [8], [13], [21]). The proof given in [13] is due to a version of equivariant Morita equivalence (cf. [4], [5], [6], [24]).

In this section, we assume that two matrices A and B are elementary equivalent such that $A = CD$ and $B = DC$. We set the square matrix $Z = \begin{bmatrix} 0 & C \\ D & 0 \end{bmatrix}$ so that we see

$$Z^2 = \begin{bmatrix} CD & 0 \\ 0 & DC \end{bmatrix} = \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix}.$$

Let $\mathcal{O}_Z$ be the Cuntz–Krieger algebra for the matrix Z . Let us denote by $S_c, S_d, c \in E_C, d \in E_D$ the generating partial isometries of $\mathcal{O}_Z$ such that $\sum_{c \in E_C} S_c S_c^* + \sum_{d \in E_D} S_d S_d^* = 1$ and

$$S_c^* S_c = \sum_{d \in E_D} Z(c, d) S_d S_d^*, \quad S_d^* S_d = \sum_{c \in E_C} Z(d, c) S_c S_c^*$$

for $c \in E_C, d \in E_D$. We note that $S_c S_d \neq 0$ (resp. $S_d S_c \neq 0$) if and only if $\varphi_{A, CD}(a) = cd$ (resp. $\varphi_{B, DC}(b) = dc$) for some $a \in E_A$ (resp. $b \in E_B$). In this case, we identify cd (resp. dc) with a (resp. b) through the map $\varphi_{A, CD}$ (resp. $\varphi_{B, DC}$). We then write $S_{cd} = S_a$ (resp. $S_{dc} = S_b$) where $S_{cd} = S_c S_d$ (resp. $S_{dc} = S_d S_c$) belongs to $\mathcal{O}_Z$ whereas S_a (resp. S_b) belongs to $\mathcal{O}_A$ (resp. $\mathcal{O}_B$). Put $P_C = \sum_{c \in E_C} S_c S_c^*$ and $P_D = \sum_{d \in E_D} S_d S_d^*$ so that $P_C + P_D = 1$. It has been shown in [13] that

$$P_C \mathcal{O}_Z P_C = \mathcal{O}_A, \quad P_D \mathcal{O}_Z P_D = \mathcal{O}_B, \quad \mathcal{D}_Z P_C = \mathcal{D}_A, \quad \mathcal{D}_Z P_D = \mathcal{D}_B \quad (4.1)$$

and $P_C \mathcal{O}_Z P_D$ has a natural structure of $\mathcal{O}_A - \mathcal{O}_B$ imprimitivity bimodule ([26], [27]). As $\mathcal{D}_Z$ is commutative, we note that $P_C \mathcal{D}_Z P_C = \mathcal{D}_Z P_C$ and $P_D \mathcal{D}_Z P_D = \mathcal{D}_Z P_D$.

We take and fix a function $f \in C(X_A, \mathbb{Z})$. It is regarded as an element of $\mathcal{D}_A$ and hence of $\mathcal{D}_Z$ by identifying it with $f \oplus 0$ in $\mathcal{D}_A \oplus \mathcal{D}_B = \mathcal{D}_Z$. Since $\exp(2\pi\sqrt{-1}tf(f \oplus 0)) = \exp(2\pi\sqrt{-1}tf) + P_D \in \mathcal{U}(\mathcal{D}_Z)$, the automorphism $\rho_t^{Z, f} \in \text{Aut}(\mathcal{O}_Z)$ for $t \in \mathbb{R}/\mathbb{Z}$ satisfies

$$\rho_t^{Z, f}(S_c) = \exp(2\pi\sqrt{-1}tf) S_c \quad \text{for } c \in E_C, \quad \rho_t^{Z, f}(S_d) = S_d \quad \text{for } d \in E_D. \quad (4.2)$$

For $a \in E_A, b \in E_B$ such that $\varphi_{A, CD}(a) = cd, \varphi_{B, DC}(b) = dc$, we then have

$$\begin{aligned} \rho_t^{Z, f}(S_c S_d) &= \rho_t^{Z, f}(S_c) \rho_t^{Z, f}(S_d) = \exp(2\pi\sqrt{-1}tf) S_c S_d = \rho_t^{A, f}(S_a), \\ \rho_t^{Z, f}(S_d S_c) &= \rho_t^{Z, f}(S_d) \rho_t^{Z, f}(S_c) = S_d \exp(2\pi\sqrt{-1}tf) S_c = S_d \exp(2\pi\sqrt{-1}tf) S_d^* S_b. \end{aligned}$$

Put $\phi(f) = \sum_{d \in E_D} S_d f S_d^* \in \mathcal{D}_Z$. Since $P_D \phi(f) P_D = \phi(f)$, we see that $\phi(f) \in \mathcal{D}_B$ and hence $\phi(f) \in C(X_B, \mathbb{Z})$ such that

$$\sum_{d \in E_D} S_d \exp(2\pi\sqrt{-1}tf) S_d^* = \exp(2\pi\sqrt{-1}t\phi(f)) \in \mathcal{U}(\mathcal{D}_B).$$

Therefore we see

$$\rho_t^{Z, f}(S_d S_c) = \exp(2\pi\sqrt{-1}t\phi(f)) S_b = \rho_t^{B, \phi(f)}(S_b).$$

We have thus proved the following lemma.

Lemma 4.1. *Suppose that $A = CD$ and $B = DC$. For $f \in C(X_A, \mathbb{Z})$, there exists $\phi(f) \in C(X_B, \mathbb{Z})$ such that*

$$\rho_t^{Z,f}(S_c S_d) = \rho_t^{A,f}(S_a), \quad \rho_t^{Z,f}(S_d S_c) = \rho_t^{B,\phi(f)}(S_b), \quad t \in \mathbb{R}/\mathbb{Z} \quad (4.3)$$

where $a \in E_A, b \in E_B$ and $c \in E_C, d \in E_D$ are satisfying $\varphi_{A,CD}(a) = cd, \varphi_{B,DC}(b) = dc$.

We note that the following symmetric equations to (4.3)

$$\rho_t^{Z,g}(S_d S_c) = \rho_t^{B,g}(S_b), \quad \rho_t^{Z,g}(S_c S_d) = \rho_t^{A,\psi(g)}(S_a), \quad t \in \mathbb{R}/\mathbb{Z} \quad (4.4)$$

for $g \in C(X_B, \mathbb{Z})$ and $\psi(g) = \sum_{c \in E_C} S_c g S_c^* \in C(X_A, \mathbb{Z})$ hold. We then have the following lemma.

Lemma 4.2. *Suppose that $A = CD$ and $B = DC$. The homomorphisms $\phi : C(X_A, \mathbb{Z}) \rightarrow C(X_B, \mathbb{Z})$ and $\psi : C(X_B, \mathbb{Z}) \rightarrow C(X_A, \mathbb{Z})$ defined by*

$$\phi(f) = \sum_{d \in E_D} S_d f S_d^*, \quad \psi(g) = \sum_{c \in E_C} S_c g S_c^*$$

satisfy the following equalities

$$(\psi \circ \phi)(f) = f \circ \sigma_A, \quad (\phi \circ \psi)(g) = g \circ \sigma_B \quad (4.5)$$

for $f \in C(X_A, \mathbb{Z})$ and $g \in C(X_B, \mathbb{Z})$. Hence they induce isomorphisms $\bar{\phi} : H^A \rightarrow H^B$ and $\bar{\psi} : H^B \rightarrow H^A$ of ordered abelian groups in a natural way.

Proof. We have

$$(\psi \circ \phi)(f) = \sum_{c \in E_C} S_c \left(\sum_{d \in E_D} S_d f S_d^* \right) S_c^* = \sum_{a \in E_A} S_a f S_a^* = f \circ \sigma_A \quad \text{for } f \in C(X_A, \mathbb{Z})$$

and similarly $(\phi \circ \psi)(g) = g \circ \sigma_B$ for $g \in C(X_B, \mathbb{Z})$. It is easy to see that the equality $\phi(f - f \circ \sigma_A) = \phi(f) - \phi(f) \circ \sigma_B$ holds so that $\phi : C(X_A, \mathbb{Z}) \rightarrow C(X_B, \mathbb{Z})$ induces an isomorphism written $\bar{\phi} : H^A \rightarrow H^B$ of ordered abelian groups whose inverse is $\bar{\psi} : H^B \rightarrow H^A$ induced by $\psi : C(X_B, \mathbb{Z}) \rightarrow C(X_A, \mathbb{Z})$. $\square$

We can prove the following proposition.

Proposition 4.3. *Suppose that $A = CD$ and $B = DC$. Then there exists an isomorphism $\Phi : \mathcal{O}_A \otimes \mathcal{K} \rightarrow \mathcal{O}_B \otimes \mathcal{K}$ satisfying $\Phi(\mathcal{D}_A \otimes \mathcal{C}) = \mathcal{D}_B \otimes \mathcal{C}$ such that for each function $f \in C(X_A, \mathbb{Z})$ there exists a unitary one-cocycle $u_t^f \in \mathcal{U}(M(\mathcal{O}_A \otimes \mathcal{K}))$ relative to $\rho^{A,f} \otimes \text{id}$ such that*

$$\Phi \circ \text{Ad}(u_t^f) \circ (\rho_t^{A,f} \otimes \text{id}) = (\rho_t^{B,\phi(f)} \otimes \text{id}) \circ \Phi \quad \text{for } t \in \mathbb{R}/\mathbb{Z}.$$

Proof. By [13, Proposition 4.1], there exist partial isometries $v_A, v_B \in M(\mathcal{O}_Z \otimes \mathcal{K})$ such that

$$v_A^* v_A = v_B^* v_B = 1, \quad v_A v_A^* = P_C \otimes 1, \quad v_B v_B^* = P_D \otimes 1 \quad (4.6)$$

and

$$\text{Ad}(v_A^*) : \mathcal{O}_A \otimes \mathcal{K} \rightarrow \mathcal{O}_Z \otimes \mathcal{K} \quad \text{and} \quad \text{Ad}(v_B^*) : \mathcal{O}_B \otimes \mathcal{K} \rightarrow \mathcal{O}_Z \otimes \mathcal{K} \quad (4.7)$$

are isomorphisms satisfying

$$\text{Ad}(v_A^*)(\mathcal{D}_A \otimes \mathcal{C}) = \mathcal{D}_Z \otimes \mathcal{C} \quad \text{and} \quad \text{Ad}(v_B^*)(\mathcal{D}_B \otimes \mathcal{C}) = \mathcal{D}_Z \otimes \mathcal{C}.$$

Put $w = v_B v_A^* \in M(\mathcal{O}_Z \otimes \mathcal{K})$ and $\Phi = \text{Ad}(w) : \mathcal{O}_A \otimes \mathcal{K} \rightarrow \mathcal{O}_B \otimes \mathcal{K}$ which satisfies $\Phi(\mathcal{D}_A \otimes \mathcal{C}) = \mathcal{D}_B \otimes \mathcal{C}$. For $f \in C(X_A, \mathbb{Z})$, we define $u_t^f \in \mathcal{U}(M(\mathcal{O}_A \otimes \mathcal{K}))$ by setting $u_t^f = w^*(\rho_t^{Z,f} \otimes \text{id})(w)$, where $\rho_t^{Z,f} \otimes \text{id}$ is extended to an automorphism of $M(\mathcal{O}_Z \otimes \mathcal{K})$. As $(P_C \otimes 1)v_A = v_A$, we have

$$u_t^f = (P_C \otimes 1)w^*(\rho_t^{Z,f} \otimes \text{id})(w)(P_C \otimes 1)$$

so that $u_t^f \in (P_C \otimes 1)M(\mathcal{O}_Z \otimes \mathcal{K})(P_C \otimes 1)$. Since $(P_C \otimes 1)M(\mathcal{O}_Z \otimes \mathcal{K})(P_C \otimes 1) = M(\mathcal{O}_A \otimes \mathcal{K})$, we have $u_t^f \in M(\mathcal{O}_A \otimes \mathcal{K})$. As $\rho_t^{A,f}(T) = \rho_t^{Z,f}(T)$ for $T \in \mathcal{O}_A$, it then follows that for $T \otimes K \in \mathcal{O}_A \otimes \mathcal{K}$

$$\begin{aligned} [\text{Ad}(u_t^f) \circ (\rho_t^{A,f} \otimes \text{id})](T \otimes K) &= u_t^f(\rho_t^{A,f}(T) \otimes K)(u_t^f)^* \\ &= w^*(\rho_t^{Z,f} \otimes \text{id})(w)(\rho_t^{Z,f}(T) \otimes K)(w^*(\rho_t^{Z,f} \otimes \text{id})(w))^* \\ &= w^*(\rho_t^{Z,f} \otimes \text{id})(w(T \otimes K)w^*)w. \end{aligned}$$

Since $w(T \otimes K)w^* \in \mathcal{O}_B \otimes \mathcal{K}$ and $\rho_t^{Z,f}(R) = \rho_t^{B,\phi(f)}(R)$ for $R \in \mathcal{O}_B$, we have

$$\begin{aligned} w^*(\rho_t^{Z,f} \otimes \text{id})(w(T \otimes K)w^*)w &= w^*(\rho_t^{B,\phi(f)} \otimes \text{id})(w(T \otimes K)w^*)w \\ &= [\Phi^{-1} \circ (\rho_t^{B,\phi(f)} \otimes \text{id}) \circ \Phi](T \otimes K). \end{aligned}$$

Hence we obtain

$$\text{Ad}(u_t^f) \circ (\rho_t^{A,f} \otimes \text{id}) = \Phi^{-1} \circ (\rho_t^{B,\phi(f)} \otimes \text{id}) \circ \Phi.$$

□

Corollary 4.4. *Suppose that A and B are strong shift equivalent. Then there exist an isomorphism $\Phi : \mathcal{O}_A \otimes \mathcal{K} \rightarrow \mathcal{O}_B \otimes \mathcal{K}$ satisfying $\Phi(\mathcal{D}_A \otimes \mathcal{C}) = \mathcal{D}_B \otimes \mathcal{C}$ and a homomorphism $\phi : C(X_A, \mathbb{Z}) \rightarrow C(X_B, \mathbb{Z})$ of ordered groups which induces an isomorphism between H^A and H^B of ordered groups such that for each function $f \in C(X_A, \mathbb{Z})$ there exists a unitary one-cocycle $u_t^f \in \mathcal{U}(M(\mathcal{O}_A \otimes \mathcal{K}))$ relative to $\rho^{A,f} \otimes \text{id}$ satisfying*

$$\Phi \circ \text{Ad}(u_t^f) \circ (\rho_t^{A,f} \otimes \text{id}) = (\rho_t^{B,\phi(f)} \otimes \text{id}) \circ \Phi \quad \text{for } t \in \mathbb{R}/\mathbb{Z}.$$

We note that the above isomorphism from H^A to H^B induced by ϕ sends the class $[1_A]$ of the constant 1 function on X_A to the class $[1_B]$ of the constant 1 function on X_B .

5 Flow equivalence and circle actions

For an $N \times N$ matrix $A = [A(i, j)]_{i,j=1}^N$ with entries in $\{0, 1\}$, put

$$\tilde{A} = \begin{bmatrix} 0 & A(1, 1) & \cdots & A(1, N) \\ 1 & 0 & \cdots & 0 \\ 0 & A(2, 1) & \cdots & A(2, N) \\ \vdots & \vdots & & \vdots \\ 0 & A(N, 1) & \cdots & A(N, N) \end{bmatrix}, \quad (5.1)$$

which is called the expansion of A at the vertex 1. Let $\{0, 1, \dots, N\}$ be the set of symbols for the topological Markov shifts $(X_{\tilde{A}}, \sigma_{\tilde{A}})$ for the matrix $\tilde{A}$. Let us denote by $\tilde{S}_0, \tilde{S}_1, \dots, \tilde{S}_N$ the canonical generating partial isometries of the Cuntz–Krieger algebra $\mathcal{O}_{\tilde{A}}$ which satisfy $\sum_{j=0}^N \tilde{S}_j \tilde{S}_j^* = 1, \tilde{S}_i^* \tilde{S}_i = \sum_{j=0}^N \tilde{A}(i, j) \tilde{S}_j \tilde{S}_j^*$ for $i = 0, 1, \dots, N$. Put $P = \sum_{i=1}^N \tilde{S}_i \tilde{S}_i^*$. We note that the identities

$$\tilde{S}_1^* P \tilde{S}_1 = \tilde{S}_1^* \tilde{S}_1 = \tilde{S}_0 \tilde{S}_0^*, \quad P + \tilde{S}_0 \tilde{S}_0^* = P + \tilde{S}_1^* P \tilde{S}_1 = 1 \quad (5.2)$$

hold. We decompose the set $\mathbb{N}$ of natural numbers such as $\mathbb{N} = \cup_{j=1}^{\infty} \mathbb{N}_j$ and $\mathbb{N}_j = \cup_{k=0}^{\infty} \mathbb{N}_{j_k}$ as disjoint unions such that $\mathbb{N}_{j_k}$ is an infinite set for each k, j . We take and fix a canonical system $e_{i,j}, i, j \in \mathbb{N}$ of matrix units which generates the algebra $\mathcal{K}$ such that $e_{i,i}, i \in \mathbb{N}$ are projections of rank one in $\mathcal{C}$. We put the projections $f_j = \sum_{i \in \mathbb{N}_j} e_{i,i}, f_{j_k} = \sum_{i \in \mathbb{N}_{j_k}} e_{i,i}$. Take a partial isometry $s_{j_k,j}$ such that $s_{j_k,j}^* s_{j_k,j} = f_j, s_{j_k,j} s_{j_k,j}^* = f_{j_k}$.

Lemma 5.1. *There exists a partial isometry $w_n \in M(\mathcal{O}_{\tilde{A}} \otimes \mathcal{K})$, $n = 1, 2, \dots$ such that*

$$\begin{aligned} w_n^* w_n &= 1_{\tilde{A}} \otimes f_n, & w_n w_n^* &\leq P \otimes f_n, \\ w_n^* (\mathcal{D}_{\tilde{A}} P \otimes \mathcal{C}) w_n &\subset \mathcal{D}_{\tilde{A}} \otimes \mathcal{C}, & w_n (\mathcal{D}_{\tilde{A}} \otimes \mathcal{C}) w_n^* &\subset \mathcal{D}_{\tilde{A}} P \otimes \mathcal{C}. \end{aligned}$$

Proof. Put $w_n = P \otimes s_{n_0,n} + P \tilde{S}_1 \otimes s_{n_1,n}$. It then follows that

$$w_n^* w_n = P \otimes f_n + \tilde{S}_1^* P \tilde{S}_1 \otimes f_n = (P + \tilde{S}_1^* P \tilde{S}_1) \otimes f_n = 1 \otimes f_n.$$

We note that $P \tilde{S}_1 = \tilde{S}_1$ and hence $P \tilde{S}_1 P = \tilde{S}_1 P = 0$ so that

$$w_n w_n^* = P \otimes f_{n_0} + P \tilde{S}_1 \tilde{S}_1^* P \otimes f_{n_1} \leq P \otimes (f_{n_0} + f_{n_1}) \leq P \otimes f_n.$$

For $F \otimes L \in \mathcal{D}_{\tilde{A}} \otimes \mathcal{C}$ with $F \in \mathcal{D}_{\tilde{A}}$ and $L \in \mathcal{C}$, we have $P \tilde{S}_1 F P = P F \tilde{S}_1^* P = 0$ so that

$$\begin{aligned} w_n (F \otimes L) w_n^* &= P F P \otimes s_{n_0,n} L s_{n_0,n}^* + P \tilde{S}_1 F \tilde{S}_1^* P \otimes s_{n_1,n} L s_{n_1,n}^*, \\ w_n^* (P \otimes 1) (F \otimes L) (P \otimes 1) w_n &= P F P \otimes s_{n_0,n}^* L s_{n_0,n} + \tilde{S}_1^* P F P \tilde{S}_1 \otimes s_{n_1,n}^* L s_{n_1,n} \end{aligned}$$

so that $w_n (\mathcal{D}_{\tilde{A}} \otimes \mathcal{C}) w_n^* \subset \mathcal{D}_{\tilde{A}} P \otimes \mathcal{C}$ and $w_n^* (\mathcal{D}_{\tilde{A}} P \otimes \mathcal{C}) w_n \subset \mathcal{D}_{\tilde{A}} \otimes \mathcal{C}$. $\square$

Let us denote by $f_{n,m}$ a partial isometry satisfying $f_{n,m}^* f_{n,m} = f_m, f_{n,m} f_{n,m}^* = f_n$. We set $v_1 = w_1, v_2 = (P \otimes f_1 - v_1 v_1^*) (P \otimes f_{1,2})$ and

$$v_{2n-1} = w_n (1 \otimes f_n - v_{2n-2}^* v_{2n-2}), \quad v_{2n} = (P \otimes f_n - v_{2n-1} v_{2n-1}^*) (P \otimes f_{n,n+1}) \quad \text{for } 1 < n \in \mathbb{N}.$$

We know that $\sum_{n=1}^{\infty} v_n$ converges to a partial isometry v in the multiplier algebra $M(\mathcal{O}_{\tilde{A}} \otimes \mathcal{K})$ under the strict topology. By using the above lemma, we then obtain the following lemma in a similar fashion to an argument giving in the proofs of [3, Lemma 2.5] and [13, Proposition 4.1].

Lemma 5.2. *The partial isometry $v \in M(\mathcal{O}_{\tilde{A}} \otimes \mathcal{K})$ satisfy*

$$\begin{aligned} v^* v &= 1_{\tilde{A}} \otimes 1, & v v^* &= P \otimes 1, \\ v^* (\mathcal{D}_{\tilde{A}} P \otimes \mathcal{C}) v &= \mathcal{D}_{\tilde{A}} \otimes \mathcal{C}, & v (\mathcal{D}_{\tilde{A}} \otimes \mathcal{C}) v^* &= \mathcal{D}_{\tilde{A}} P \otimes \mathcal{C}. \end{aligned}$$

We thus have

Proposition 5.3. *Let $\tilde{A}$ be the matrix defined in (5.1) for a matrix A . Then there exists an isomorphism $\tilde{\Phi} : \mathcal{O}_{\tilde{A}} \otimes \mathcal{K} \rightarrow P\mathcal{O}_{\tilde{A}}P \otimes \mathcal{K}$ satisfying $\tilde{\Phi}(\mathcal{D}_{\tilde{A}} \otimes \mathcal{C}) = \mathcal{D}_{\tilde{A}}P \otimes \mathcal{C}$ such that for each function $\tilde{f} \in C(X_{\tilde{A}}, \mathbb{Z})$ there exists a unitary one-cocycle $v_t^{\tilde{f}} \in \mathcal{U}(M(\mathcal{O}_{\tilde{A}} \otimes \mathcal{K}))$ relative to $\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id}$ such that*

$$\tilde{\Phi} \circ \text{Ad}(v_t^{\tilde{f}}) \circ (\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id}) = (\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id}) \circ \tilde{\Phi} \quad \text{for } t \in \mathbb{R}/\mathbb{Z}.$$

Proof. Take a partial isometry $v \in M(\mathcal{O}_{\tilde{A}} \otimes \mathcal{K})$ as in the above lemma. Define $\tilde{\Phi} : \mathcal{O}_{\tilde{A}} \otimes \mathcal{K} \rightarrow P\mathcal{O}_{\tilde{A}}P \otimes \mathcal{K}$ by setting $\tilde{\Phi}(T \otimes K) = v(T \otimes K)v^*$ for $T \in \mathcal{O}_{\tilde{A}}$ and $K \in \mathcal{K}$. We set $v_t^{\tilde{f}} = v^*(\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id})(v)$ for $t \in \mathbb{R}/\mathbb{Z}$. It then follows that

$$\begin{aligned} (\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id}) \circ \tilde{\Phi}(T \otimes K) &= (\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id})(v(T \otimes K)v^*) \\ &= (\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id})((P \otimes 1)v(T \otimes K)v^*(P \otimes 1)) \\ &= (P \otimes 1)(\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id})(v(T \otimes K)v^*)(P \otimes 1) \\ &= vv^*(\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id})(v)(\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id})(T \otimes K)(\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id})(v^*)vv^* \\ &= \tilde{\Phi} \circ \text{Ad}(v_t^{\tilde{f}}) \circ (\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id})(T \otimes K). \end{aligned}$$

Hence we have

$$\tilde{\Phi} \circ \text{Ad}(v_t^{\tilde{f}}) \circ (\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id}) = (\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id}) \circ \tilde{\Phi} \quad \text{for } t \in \mathbb{R}/\mathbb{Z}.$$

□

Let $\xi : X_A \rightarrow X_{\tilde{A}}$ be the continuous map defined by substituting the symbol 1 by the word $(1, 0)$ in X_A such as

$$(1, 2, 1, 1, 2, 1, 1, 1, 2, \dots) \in X_A \rightarrow (1, 0, 2, 1, 0, 1, 0, 2, 1, 0, 1, 0, 1, 0, 2, \dots) \in X_{\tilde{A}}. \quad (5.3)$$

We define continuous functions $k_1, l_1 \in C(X_A, \mathbb{Z}_+)$ by setting

$$k_1(x) = 0, \quad l_1(x) = \begin{cases} 2 & \text{if } x_1 = 1, \\ 1 & \text{otherwise} \end{cases}$$

for $x = (x_n)_{n \in \mathbb{N}} \in X_A$ so that $\xi : X_A \rightarrow X_{\tilde{A}}$ satisfies the identity

$$\sigma_{\tilde{A}}^{k_1(x)}(\xi(\sigma_A(x))) = \sigma_{\tilde{A}}^{l_1(x)}(\xi(x)) \quad \text{for } x \in X_A. \quad (5.4)$$

Hence $\xi : X_A \rightarrow X_{\tilde{A}}$ is a continuous orbit map in the sense of [16]. We define $\Psi_{\xi}(\tilde{f}) \in C(X_A, \mathbb{Z})$ for $\tilde{f} \in C(X_{\tilde{A}}, \mathbb{Z})$ in the same formula as (1.5) so that

$$\Psi_{\xi}(\tilde{f})(x) = \begin{cases} \tilde{f}(\xi(x)) + \tilde{f}(\sigma_{\tilde{A}}(\xi(x))) & \text{if } x_1 = 1, \\ \tilde{f}(\xi(x)) & \text{otherwise} \end{cases}$$

for $x = (x_n)_{n \in \mathbb{N}} \in X_A$. As in the proof of [8, 4.1 Theorem], the partial isometries $\tilde{S}_1\tilde{S}_0, \tilde{S}_2, \dots, \tilde{S}_N$ satisfy the relations (1.2) so that we may regard them as the generating partial isometries $S_1, S_2, \dots, S_N$ of $\mathcal{O}_A$ and put $S_1 = \tilde{S}_1\tilde{S}_0, S_2 = \tilde{S}_2, \dots, S_N = \tilde{S}_N$.

Lemma 5.4 (cf. [8, 4.1 Theorem]). *The identities*

$$P\tilde{S}_1\tilde{S}_0P = S_1, \quad P\tilde{S}_iP = S_i \quad \text{for } i = 2, \dots, N \quad (5.5)$$

hold and the correspondence $\Phi^0(P\tilde{S}_1\tilde{S}_0P) = S_1, \Phi^0(P\tilde{S}_iP) = S_i, i = 2, \dots, N$ gives rise to an isomorphism $\Phi^0 : P\mathcal{O}_{\tilde{A}}P \rightarrow \mathcal{O}_A$ such that $\Phi^0(\mathcal{D}_{\tilde{A}}P) = \mathcal{D}_A$ and

$$\Phi^0 \circ \rho_t^{\tilde{A}, \tilde{f}} = \rho_t^{A, \Psi_\xi(\tilde{f})} \circ \Phi^0 \quad \text{for } \tilde{f} \in C(X_{\tilde{A}}, \mathbb{Z}), t \in \mathbb{R}/\mathbb{Z}.$$

Proof. Since $P\tilde{S}_1 = \tilde{S}_1$ and $\tilde{S}_0P = \tilde{S}_0$, we have $P\tilde{S}_1\tilde{S}_0P = S_1$. It is easy to see that $P\tilde{S}_iP = S_i$ for $i = 2, \dots, N$. As in the proof of [8, 4.1 Theorem], the correspondence $\Phi^0 : P\mathcal{O}_{\tilde{A}}P \rightarrow \mathcal{O}_A$ above is shown to be an isomorphism satisfying $\Phi^0(\mathcal{D}_{\tilde{A}}P) = \mathcal{D}_A$. We note that $U_t(\tilde{f}^2)\tilde{S}_1\tilde{S}_0 = U_t(\Psi_\xi(\tilde{f}))S_1$ which implies that $\Phi^0(U_t(\tilde{f}^2)\tilde{S}_1\tilde{S}_0) = U_t(\Psi_\xi(\tilde{f}))S_1$. It then follows that

$$\begin{aligned} \Phi^0 \circ \rho_t^{\tilde{A}, \tilde{f}}(P\tilde{S}_1\tilde{S}_0P) &= \Phi^0(\rho_t^{\tilde{A}, \tilde{f}}(\tilde{S}_1\tilde{S}_0)) = \Phi^0(U_t(\tilde{f}^2)\tilde{S}_1\tilde{S}_0) \\ &= U_t(\Psi_\xi(\tilde{f}))S_1 = \rho_t^{A, \Psi_\xi(\tilde{f})}(\Phi^0(P\tilde{S}_1\tilde{S}_0P)). \end{aligned}$$

Similarly

$$\Phi^0 \circ \rho_t^{\tilde{A}, \tilde{f}}(P\tilde{S}_iP) = \rho_t^{A, \Psi_\xi(\tilde{f})}(\Phi^0(P\tilde{S}_iP)), \quad i = 2, \dots, N.$$

Hence we obtain that $\Phi^0 \circ \rho_t^{\tilde{A}, \tilde{f}} = \rho_t^{A, \Psi_\xi(\tilde{f})} \circ \Phi^0$. $\square$

Let $\eta : X_{\tilde{A}} \rightarrow X_A$ be the continuous map defined by deleting the symbol 0 in elements of $X_{\tilde{A}}$ such as

$$\begin{aligned} (1, 0, 2, 1, 0, 1, 0, 2, 1, 0, 1, 0, 1, 0, 2, \dots) &\in X_{\tilde{A}} \rightarrow (1, 2, 1, 1, 2, 1, 1, 1, 2, \dots) \in X_A, \\ (0, 2, 1, 0, 1, 0, 2, 1, 0, 1, 0, 1, 0, 2, 1, \dots) &\in X_{\tilde{A}} \rightarrow (2, 1, 1, 2, 1, 1, 1, 2, 1, \dots) \in X_A. \end{aligned}$$

We define continuous functions $\tilde{k}_1, \tilde{l}_1 \in C(X_{\tilde{A}}, \mathbb{Z}_+)$ by setting

$$\tilde{k}_1(\tilde{x}) = 0, \quad \tilde{l}_1(\tilde{x}) = \begin{cases} 0 & \text{if } \tilde{x}_1 = 0, \\ 1 & \text{otherwise} \end{cases}$$

for $\tilde{x} = (\tilde{x}_n)_{n \in \mathbb{N}} \in X_{\tilde{A}}$ so that $\eta : X_{\tilde{A}} \rightarrow X_A$ satisfies the identity

$$\sigma_A^{\tilde{k}_1(\tilde{x})}(\eta(\sigma_{\tilde{A}}(\tilde{x}))) = \sigma_A^{\tilde{l}_1(\tilde{x})}(\eta(\tilde{x})) \quad \text{for } \tilde{x} \in X_{\tilde{A}}. \quad (5.6)$$

Hence $\eta : X_{\tilde{A}} \rightarrow X_A$ is a continuous orbit map in the sense of [16]. We define $\Psi_\eta(f) \in C(X_{\tilde{A}}, \mathbb{Z})$ for $f \in C(X_A, \mathbb{Z})$ in the same formula as (1.5) so that

$$\Psi_\eta(f)(\tilde{x}) = \begin{cases} 0 & \text{if } \tilde{x}_1 = 0, \\ f(\eta(\tilde{x})) & \text{otherwise} \end{cases}$$

for $\tilde{x} = (\tilde{x}_n)_{n \in \mathbb{N}} \in X_{\tilde{A}}$. We then have

Lemma 5.5. *The homomorphisms $\Psi_\eta : C(X_A, \mathbb{Z}) \rightarrow C(X_{\tilde{A}}, \mathbb{Z})$ and $\Psi_\xi : C(X_{\tilde{A}}, \mathbb{Z}) \rightarrow C(X_A, \mathbb{Z})$ induce isomorphisms of ordered abelian groups $\tilde{\Psi}_\eta : H^A \rightarrow H^{\tilde{A}}$ and $\tilde{\Psi}_\xi : H^{\tilde{A}} \rightarrow H^A$, respectively such that they are inverses to each other.*

Proof. Both the maps $\eta : X_{\tilde{A}} \rightarrow X_A$ and $\xi : X_A \rightarrow X_{\tilde{A}}$ are continuous orbit maps so that the homomorphisms $\Psi_\eta : C(X_A, \mathbb{Z}) \rightarrow C(X_{\tilde{A}}, \mathbb{Z})$ and $\Psi_\xi : C(X_{\tilde{A}}, \mathbb{Z}) \rightarrow C(X_A, \mathbb{Z})$ induce homomorphisms of ordered abelian groups $\bar{\Psi}_\eta : H^A \rightarrow H^{\tilde{A}}$ and $\bar{\Psi}_\xi : H^{\tilde{A}} \rightarrow H^A$, respectively ([16]). It suffices to prove that they are inverses to each other. It is easy to see that

$$\begin{aligned} \eta \circ \xi(x) &= x \quad \text{for } x \in X_A, \\ \xi \circ \eta(\tilde{x}) &= \begin{cases} \sigma_{\tilde{A}}(\tilde{x}) & \text{if } \tilde{x}_1 = 0, \\ \tilde{x} & \text{otherwise} \end{cases} \quad \text{for } \tilde{x} = (\tilde{x}_n)_{n \in \mathbb{N}} \in X_{\tilde{A}}. \end{aligned}$$

As in [16, Proposition 3.4], $\eta \rightarrow \Psi_\eta$ is contravariantly functorial, so that we have

$$\begin{aligned} (\Psi_\xi \circ \Psi_\eta)(f) &= \Psi_{\eta \circ \xi}(f) = f \quad \text{for } f \in C(X_A, \mathbb{Z}), \\ (\Psi_\eta \circ \Psi_\xi)(\tilde{f})(\tilde{x}) &= \Psi_{\xi \circ \eta}(\tilde{f})(\tilde{x}) = \begin{cases} 0 & \text{if } \tilde{x}_1 = 0, \\ \tilde{f}(\tilde{x}) + \tilde{f}(\sigma_{\tilde{A}}(\tilde{x})) & \text{if } \tilde{x}_1 = 1, \\ \tilde{f}(\tilde{x}) & \text{otherwise} \end{cases} \end{aligned}$$

for $\tilde{x} = (\tilde{x}_n)_{n \in \mathbb{N}} \in X_{\tilde{A}}$. We set

$$\tilde{f}_0(\tilde{x}) = \begin{cases} \tilde{f}(\tilde{x}) & \text{if } \tilde{x}_1 = 0, \\ 0 & \text{otherwise.} \end{cases}$$

It is straightforward to see that

$$(\Psi_\eta \circ \Psi_\xi)(\tilde{f})(\tilde{x}) - \tilde{f}(\tilde{x}) = \tilde{f}_0(\sigma_{\tilde{A}}(\tilde{x})) - \tilde{f}_0(\tilde{x}) \quad \text{for } \tilde{x} \in X_{\tilde{A}}$$

so that $(\Psi_\eta \circ \Psi_\xi)(\tilde{f}) - \tilde{f} = \tilde{f}_0 \circ \sigma_{\tilde{A}} - \tilde{f}_0$ and hence $[(\Psi_\eta \circ \Psi_\xi)(\tilde{f})] = [\tilde{f}]$ for all $\tilde{f} \in C(X_{\tilde{A}}, \mathbb{Z})$. $\square$

By combining Proposition 5.3 with Lemma 5.4 and Lemma 5.5, we have the following proposition.

Proposition 5.6. *Let $\tilde{A}$ be the matrix defined by (5.1) for a matrix A . Then there exist an isomorphism $\Phi : \mathcal{O}_{\tilde{A}} \otimes \mathcal{K} \rightarrow \mathcal{O}_A \otimes \mathcal{K}$ satisfying $\Phi(\mathcal{D}_{\tilde{A}} \otimes \mathcal{C}) = \mathcal{D}_A \otimes \mathcal{C}$, a homomorphism $\Psi_\xi : C(X_{\tilde{A}}, \mathbb{Z}) \rightarrow C(X_A, \mathbb{Z})$ inducing an isomorphism $\bar{\Psi}_\xi : H^{\tilde{A}} \rightarrow H^A$ of ordered abelian groups, and a unitary one-cocycle $v_t^{\tilde{f}} \in \mathcal{U}(M(\mathcal{O}_{\tilde{A}} \otimes \mathcal{K}))$ relative to $\rho^{\tilde{A}, \tilde{f}} \otimes \text{id}$ for each function $\tilde{f} \in C(X_{\tilde{A}}, \mathbb{Z})$ such that*

$$\Phi \circ \text{Ad}(v_t^{\tilde{f}}) \circ (\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id}) = (\rho_t^{A, \Psi_\xi(\tilde{f})} \otimes \text{id}) \circ \Phi \quad \text{for } t \in \mathbb{R}/\mathbb{Z}.$$

Proof. By Lemma 5.4 there exists an isomorphism

$$\Phi^0 : P\mathcal{O}_{\tilde{A}}P \rightarrow \mathcal{O}_A \quad \text{such that} \quad \Phi^0(\mathcal{D}_{\tilde{A}}P) = \mathcal{D}_A$$

and for $\tilde{f} \in C(X_{\tilde{A}}, \mathbb{Z})$

$$\Phi^0 \circ \rho_t^{\tilde{A}, \tilde{f}} = \rho_t^{A, \Psi_\xi(\tilde{f})} \circ \Phi^0.$$

By Proposition 5.3 there exist an isomorphism

$$\tilde{\Phi} : \mathcal{O}_{\tilde{A}} \otimes \mathcal{K} \rightarrow P\mathcal{O}_{\tilde{A}}P \otimes \mathcal{K} \quad \text{such that} \quad \tilde{\Phi}(\mathcal{D}_{\tilde{A}} \otimes \mathcal{C}) = \mathcal{D}_{\tilde{A}}P \otimes \mathcal{C}$$

and a unitary one-cocycle $v_t^{\tilde{f}} \in \mathcal{U}(M(\mathcal{O}_{\tilde{A}} \otimes \mathcal{K}))$ relative to $\rho^{\tilde{A}, \tilde{f}} \otimes \text{id}$ such that

$$\tilde{\Phi} \circ \text{Ad}(v_t^{\tilde{f}}) \circ (\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id}) = (\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id}) \circ \tilde{\Phi}.$$

We set

$$\Phi = (\Phi^0 \otimes \text{id}) \circ \tilde{\Phi} : \mathcal{O}_{\tilde{A}} \otimes \mathcal{K} \xrightarrow{\tilde{\Phi}} P\mathcal{O}_{\tilde{A}}P \otimes \mathcal{K} \xrightarrow{\Phi^0 \otimes \text{id}} \mathcal{O}_A \otimes \mathcal{K}$$

so that $\Phi(\mathcal{D}_{\tilde{A}} \otimes \mathcal{C}) = \mathcal{D}_A \otimes \mathcal{C}$ and for $\tilde{f} \in C(X_{\tilde{A}}, \mathbb{Z})$, we have

$$\begin{aligned} \Phi \circ \text{Ad}(v_t^{\tilde{f}}) \circ (\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id}) &= (\Phi^0 \otimes \text{id}) \circ \tilde{\Phi} \circ \text{Ad}(v_t^{\tilde{f}}) \circ (\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id}) \\ &= (\Phi^0 \otimes \text{id}) \circ (\rho_t^{\tilde{A}, \tilde{f}} \otimes \text{id}) \circ \tilde{\Phi} \\ &= [(\Phi^0 \circ \rho_t^{\tilde{A}, \tilde{f}}) \otimes \text{id}] \circ \tilde{\Phi} \\ &= [(\rho_t^{A, \Psi_\xi(\tilde{f})} \circ \Phi^0) \otimes \text{id}] \circ \tilde{\Phi} \\ &= (\rho_t^{A, \Psi_\xi(\tilde{f})} \otimes \text{id}) \circ \Phi. \end{aligned}$$

□

In [22], Parry–Sullivan proved that the flow equivalence relation of topological Markov shifts is generated by the expansions $A \rightarrow \tilde{A}$ and topological conjugacies. Therefore we arrive at the following theorem.

Theorem 5.7. *If two-sided topological Markov shifts $(\bar{X}_A, \bar{\sigma}_A)$ and $(\bar{X}_B, \bar{\sigma}_B)$ are flow equivalent, then there exist an isomorphism $\Phi : \mathcal{O}_A \otimes \mathcal{K} \rightarrow \mathcal{O}_B \otimes \mathcal{K}$, a homomorphism $\varphi : C(X_A, \mathbb{Z}) \rightarrow C(X_B, \mathbb{Z})$ of ordered abelian groups and a unitary one-cocycle $u_t^f \in \mathcal{U}(M(\mathcal{O}_A \otimes \mathcal{K}))$ relative to $\rho^{A, f} \otimes \text{id}$ for each function $f \in C(X_A, \mathbb{Z})$ satisfying the following conditions*

- (i) $\Phi(\mathcal{D}_A \otimes \mathcal{C}) = \mathcal{D}_B \otimes \mathcal{C}$,
- (ii) $\Phi \circ \text{Ad}(u_t^f) \circ (\rho_t^{A, f} \otimes \text{id}) = (\rho_t^{B, \varphi(f)} \otimes \text{id}) \circ \Phi$ for $f \in C(X_A, \mathbb{Z})$, $t \in \mathbb{R}/\mathbb{Z}$,
- (iii) $\varphi : C(X_A, \mathbb{Z}) \rightarrow C(X_B, \mathbb{Z})$ induces an isomorphism from H^A to H^B as ordered abelian groups.

Proof. By the Parry–Sullivan’s result, we obtain the assertion from Proposition 5.6 and Corollary 4.4. □

Recall that $\gamma_r^{A, f} \in \text{Aut}(\mathcal{O}_A)$, $r \in \mathbb{R}$ stand for the one-parameter automorphisms for $f \in C(X_A, \mathbb{R})$ defined by (3.10). We may similarly show the following result:

Proposition 5.8. *If two-sided topological Markov shifts $(\bar{X}_A, \bar{\sigma}_A)$ and $(\bar{X}_B, \bar{\sigma}_B)$ are flow equivalent, then there exists an isomorphism $\Phi : \mathcal{O}_A \otimes \mathcal{K} \rightarrow \mathcal{O}_B \otimes \mathcal{K}$ such that $\Phi(\mathcal{D}_A \otimes \mathcal{C}) = \mathcal{D}_B \otimes \mathcal{C}$ and for each function $f \in C(X_A, \mathbb{R})$ there exists a function $\tilde{f} \in C(X_B, \mathbb{R})$ and a unitary one-cocycle $u_r^f \in \mathcal{U}(M(\mathcal{O}_A \otimes \mathcal{K}))$ relative to $\gamma^{A, f} \otimes \text{id}$ such that*

$$\Phi \circ \text{Ad}(u_r^f) \circ (\gamma_r^{A, f} \otimes \text{id}) = (\gamma_r^{B, \tilde{f}} \otimes \text{id}) \circ \Phi \quad \text{for } r \in \mathbb{R}.$$

Acknowledgments: The author would like to thank the referee for his careful reading the manuscript and various suggestions. This work was supported by JSPS KAKENHI Grant Numbers 23540237 and 15K04896.

References

- [1] M. BOYLE AND D. HANDELMAN, *Orbit equivalence, flow equivalence and ordered cohomology*, Israel J. Math. **95**(1996), pp. 169–210.
- [2] R. BOWEN AND J. FRANKS, *Homology for zero-dimensional nonwandering sets*, Ann. Math. **106**(1977), pp. 73–92.
- [3] L. G. BROWN, *Stable isomorphism of hereditary subalgebras of C^* -algebras*, Pacific J. Math. **71**(1977), pp. 335–348.
- [4] L. G. BROWN, P. GREEN AND M. A. RIEFFEL, *Stable isomorphism and strong Morita equivalence of C^* -algebras*, Pacific J. Math. **71**(1977), pp. 349–363.
- [5] F. COMBES, *Crossed products and Morita equivalence*, Proc. London Math. Soc. **49**(1984), pp. 289–306.
- [6] D. CROCKER, A. KUMJIAN, I. RAEBURN AND D. P. WILLIAMS, *An equivariant Brauer group and actions of groups of C^* -algebras*, J. Funct. Anal. **146**(1997), pp. 151–184.
- [7] J. CUNTZ, *Automorphisms of certain simple C^* -algebras*, Quantum Fields-Algebras, Processes, Springer Verlag, Wien-New York, 1980, pp. 187–196.
- [8] J. CUNTZ AND W. KRIEGER, *A class of C^* -algebras and topological Markov chains*, Invent. Math. **56**(1980), pp. 251–268.
- [9] J. FRANKS, *Flow equivalence of subshifts of finite type*, Ergodic Theory Dynam. Systems **4**(1984), pp. 53–66.
- [10] B. P. KITCHENS, *Symbolic dynamics*, Springer-Verlag, Berlin, Heidelberg and New York (1998).
- [11] D. LIND AND B. MARCUS, *An introduction to symbolic dynamics and coding*, Cambridge University Press, Cambridge (1995).
- [12] K. MATSUMOTO, *On automorphisms of C^* -algebras associated with subshifts*, J. Operator Theory **44**(2000), pp. 91–112.
- [13] K. MATSUMOTO, *Strong shift equivalence of symbolic dynamical systems and Morita equivalence of C^* -algebras*, Ergodic Theory Dynam. Systems **24**(2002), pp. 199–215.
- [14] K. MATSUMOTO, *Orbit equivalence of topological Markov shifts and Cuntz–Krieger algebras*, Pacific J. Math. **246**(2010), 199–225.
- [15] K. MATSUMOTO, *Classification of Cuntz–Krieger algebras by orbit equivalence of topological Markov shifts*, Proc. Amer. Math. Soc. **141**(2013), pp. 2329–2342.

- [16] K. MATSUMOTO, *Strongly continuous orbit equivalence of one-sided topological Markov shifts*, J. Operator Theory **74**(2015), pp. 101–127.
- [17] K. MATSUMOTO AND H. MATUI, *Continuous orbit equivalence of topological Markov shifts and Cuntz–Krieger algebras*, Kyoto J. Math. **54**(2014), pp. 863–878.
- [18] K. MATSUMOTO AND H. MATUI, *Continuous orbit equivalence of topological Markov shifts and dynamical zeta functions*, preprint, arXiv:1403.0719, to appear in Ergodic Theory Dynam. Systems.
- [19] H. MATUI, *Homology and topological full groups of étale groupoids on totally disconnected spaces*, Proc. London Math. Soc. **104**(2012), pp. 27–56.
- [20] H. MATUI, *Topological full groups of one-sided shifts of finite type*, J. Reine Angew. Math. **705**(2015), pp. 35–84.
- [21] P. S. MUHLY, D. PASK AND M. TOMFORDE, *Strong shift equivalence of C^* -correspondences*, Israel J. Math. **167** (2008), pp. 315–346.
- [22] W. PARRY AND D. SULLIVAN, *A topological invariant for flows on one-dimensional spaces*, Topology **14**(1975), pp. 297–299.
- [23] Y. T. POON, *A K -theoretic invariant for dynamical systems*, Trans. Amer. Math. Soc. **311**(1989), pp. 513–533.
- [24] I. RAEBURN AND D. P. WILLIAMS, *Morita equivalence and continuous-trace C^* -algebras*, Mathematical Surveys and Monographs, vol(60) Amer. Math. Soc, (1998).
- [25] J. RENAULT, *A groupoid approach to C^* -algebras*, Lecture Notes in Math. 793, Springer-Verlag, Berlin, Heidelberg and New York (1980).
- [26] M. A. RIEFFEL, *Induced representations of C^* -algebras*, Adv. in Math. **13**(1974), pp. 176–257.
- [27] M. A. RIEFFEL, *Morita equivalence for C^* -algebras and W^* -algebras*, J. Pure Appl. Algebra **5**(1974), pp. 51–96.
- [28] M. RØRDAM, *Classification of Cuntz–Krieger algebras*, K-theory **9**(1995), pp. 31–58.
- [29] M. TOMFORDE *The Graph Algebra Problem Page : List of Open Problems*, <http://www.math.uh.edu/tomforde/GraphAlgebraProblems/ListOfProblems.html>.
- [30] R. F. WILLIAMS, *Classification of subshifts of finite type*, Ann. Math. **98**(1973), pp. 120–153. erratum, Ann. Math. **99**(1974), pp. 380–381.